



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



N-butyryl-L-Homoserine lactone

Item No. 10007898

CAS Registry No.: 67605-85-0
Formal Name: N-[(3S)-tetrahydro-2-oxo-3-furanyl]-butanamide
Synonyms: BHL, C4-HSL, PAI-2
MF: $C_8H_{13}NO_3$
FW: 171.2
Purity: $\geq 95\%$
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥ 2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

N-butyryl-L-Homoserine lactone is supplied as a crystalline solid. A stock solution may be made by dissolving the N-butyryl-L-homoserine lactone in the solvent of choice, which should be purged with an inert gas. N-butyryl-L-Homoserine lactone is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of N-butyryl-L-homoserine lactone in these solvents is approximately 30 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of N-butyryl-L-homoserine lactone can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of N-butyryl-L-homoserine lactone in PBS (pH 7.2) is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

N-butyryl-L-Homoserine lactone is a quorum-sensing signaling molecule produced by *P. aeruginosa*.² It induces expression of the virulence genes *lasB* and *rhlA* in *P. aeruginosa* when used at a concentration of $10\ \mu\text{M}$.¹ N-butyryl-L-Homoserine lactone ($50\ \mu\text{M}$) induces rhamnolipid accumulation in *P. aeruginosa* growth media.³

References

- Ikeda, T., Kajiyama, K., Kita, T., *et al.* The synthesis of optically pure enantiomers of N-acyl-homoserine lactone autoinducers and their analogues. *Chem. Lett.* **30(4)**, 314-315 (2001).
- Pearson, J.P., Passador, L., and Igleswski, B.H. A second N-acylhomoserine lactone signal produced by *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. USA.* **92(5)**, 1490-1494 (1995).
- Schooling, S.R., Charaf, U.K., Allison, D.G., *et al.* A role for rhamnolipid in biofilm dispersion. *Biofilm* **1(2)**, 91-99 (2004).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/14/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM