



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# PRODUCT INFORMATION




## sPLA<sub>2</sub> (human, recombinant Type V)

Item No. 10009563

### Overview and Properties

Synonyms:	gVPLA <sub>2</sub> , Secretory Phospholipase A <sub>2</sub> (Group V)
Source:	Recombinant protein expressed in <i>E. coli</i>
Uniprot No.:	P39877
Molecular Weight:	13.72 kDa
Storage:	-80°C (as supplied); avoid freeze/thaw cycles by aliquoting protein
Stability:	≥1 year
Purity:	≥95% estimated by SDS-PAGE
Supplied in:	50 mM Tris-HCl, pH 8.0, containing 100 mM sodium chloride, 50 mM calcium chloride, and 20% glycerol
Protein	
Concentration:	batch specific mg/ml
Activity:	batch specific U/ml
Specific Activity:	batch specific U/mg
Unit Definition:	One unit is defined as the amount of enzyme required to release 1 μmol of TNB per minute under the following condition: 25 mM Tris-HCl, pH 7.5, containing 10 mM CaCl <sub>2</sub> , 100 mM KCl, 0.3 mM Triton X-100, 1 mg/ml BSA, 500 μM 1,2-bis(heptanoylthio) Glycerophosphocholine (Item No. 62235), and 0.5 mM DTNB at 37°C.

### Images



Representative gel image shown; actual purity may vary between each batch.

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 - USA

PHONE: [800] 364-9897  
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM

# PRODUCT INFORMATION



## Description

Phospholipase A<sub>2</sub> (PLA<sub>2</sub>) catalyzes the hydrolysis of fatty acids at the sn-2 position of glycerophospholipids. PLA<sub>2</sub> (Type V) is a secretory PLA<sub>2</sub> (sPLA<sub>2</sub>) of approximately 14 kDa and is one of the isoforms in the growing list of the PLA<sub>2</sub> enzyme family.<sup>1</sup> This enzyme, rather than the sPLA<sub>2</sub> (Type II), is responsible for arachidonic acid mobilization leading to prostaglandin production in macrophages and mast cells.<sup>2-4</sup> Consistent with this role, sPLA<sub>2</sub> (Type V) is associated with the golgi apparatus, nuclear envelope, and plasma membrane in mouse bone marrow-derived mast cells.<sup>5</sup> sPLA<sub>2</sub>(Type V) has been cloned from a variety of species including human, mouse, and rat.<sup>6-8</sup> The enzyme is expressed in heart, lung, placenta, and spleen, as well as P388D1 macrophages and mast cells.<sup>2,3,8</sup>

The specific activity of Cayman's sPLA<sub>2</sub> (human, recombinant Type V) was established using 1,2-bis(heptanoylthio) Glycerophosphocholine (Item No. 62235) as the substrate. The reaction was inhibited by Thioetheramide-PC (Item No. 62750).

## References

1. Balsinde, J., Winstead, M.V., and Dennis, E.A. Phospholipase A<sub>2</sub> regulation of arachidonic acid mobilization. *FEBS Lett.* **531**(1), 2-6 (2002).
2. Balboa, M.A., Balsinde, J., Winstead, M.V., et al. Novel group V phospholipase A<sub>2</sub> involved in arachidonic acid mobilization in murine P388D<sub>1</sub> macrophages. *J. Biol. Chem.* **271**(50), 32381-32384 (1996).
3. Reddy, S.T., Winstead, M.V., Tischfield, J.A., et al. Analysis of the secretory phospholipase A<sub>2</sub> that mediates prostaglandin production in mast cells. *J. Biol. Chem.* **272**(21), 13591-13596 (1997).
4. Bingham, C.O., III, Murakami, M., Fujishima, H., et al. A heparin-sensitive phospholipase A<sub>2</sub> and prostaglandin endoperoxide synthase-2 are functionally linked in the delayed phase of prostaglandin D<sub>2</sub> generation in mouse bone marrow-derived mast cells. *J. Biol. Chem.* **271**(42), 25936-25944 (1996).
5. Bingham, C.O., III, Fijneman, R.J.A., Friend, D.S., et al. Low molecular weight group IIA and group V phospholipase A<sub>2</sub> enzymes have different intracellular locations in mouse bone marrow-derived mast cells. *J. Biol. Chem.* **274**(44), 31476-31484 (1999).
6. Chen, J., Engle, S.J., Seilhamer, J.J., et al. Cloning and recombinant expression of a novel human low molecular weight Ca<sup>2+</sup>-dependent phospholipase A<sub>2</sub>. *J. Biol. Chem.* **269**(4), 2365-2368 (1994).
7. Tischfield, J.A., Xia, Y.R., Shih, D.M., et al. Low-molecular-weight, calcium-dependent phospholipase A<sub>2</sub> genes are linked and map to homologous chromosome regions in mouse and human. *Genomics* **32**(3), 328-333 (1996).
8. Chen, J., Engle, S.J., Seilhamer, J.J., et al. Cloning, expression and partial characterization of a novel rat phospholipase A<sub>2</sub>. *Biochim. Biophys. Acta* **1215**(1-2), 115-120 (1994).

## CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 - USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

[CUSTSERV@CAYMANCHEM.COM](mailto:CUSTSERV@CAYMANCHEM.COM)  
[WWW.CAYMANCHEM.COM](http://WWW.CAYMANCHEM.COM)