



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



Lomitapide

Item No. 10009610

CAS Registry No.: 182431-12-5

Formal Name: N-(2,2,2-trifluoroethyl)-9-[4-[4'-[[[4'-(trifluoromethyl)[1,1'-biphenyl]-2-yl]carbonyl]amino]-1-piperidinyl]butyl]-9H-fluorene-9-carboxamide

Synonyms: AEGR 773, BMS 201038

MF: $C_{39}H_{37}F_6N_3O_2$

FW: 693.7


Purity: $\geq 98\%$

UV/Vis.: λ_{max} : 232, 259 nm

Supplied as: A crystalline solid

Storage: -20°C

Stability: ≥ 2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Lomitapide is supplied as a crystalline solid. A stock solution may be made by dissolving the lomitapide in the solvent of choice. Lomitapide is soluble in organic solvents such as ethanol and DMSO, which should be purged with an inert gas. The solubility of lomitapide in these solvents is approximately 10 and 5 mg/ml, respectively.

Description

Lomitapide is an inhibitor of microsomal triglyceride transfer protein (MTTP; $IC_{50} = 0.5$ nM in a triglyceride transfer assay).¹ It inhibits apolipoprotein B (ApoB) secretion in HepG2 cells ($EC_{50} = 0.8$ nM). Lomitapide inhibits triglyceride secretion in fasted rats ($ED_{50} = 0.19$ mg/kg, p.o.) and reduces total plasma cholesterol levels in hamsters ($ED_{50} = 2.4$ mg/kg). It also decreases plasma cholesterol and triglyceride levels in the Watanabe-heritable hyperlipidemic (WHHL) rabbit model of homozygous familial hypercholesterolemia when administered at a dose of 10 mg/kg. Formulations containing lomitapide have been used as an adjunct to a low-fat diet and other lipid-lowering treatments in the treatment of homozygous familial hypercholesterolemia.

Reference

1. Wetterau, J.R., Gregg, R.E., Harrity, T.W., *et al.* An MTP inhibitor that normalizes atherogenic lipoprotein levels in WHHL rabbits. *Science* **282**(5389), 751-754 (1998).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/11/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM