

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Dimethoxycurcumin

Item No. 10009986

CAS Registry No.:	160096-59-3		
Formal Name:	1,7-bis(3,4-dimethoxyphenyl)-1,6-		
	heptadiene-3,5-dione	0 0	
Synonym:	Di-O-methylcurcumin		
MF:	C ₂₃ H ₂₄ O ₆		
FW:	396.4		/
Purity:	≥90%	0	
UV/Vis.:	λ _{max} : 418 nm	Ó	
Supplied as:	A crystalline solid		
Storage:	-20°C		
Stability:	≥2 years		

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Dimethoxycurcumin is supplied as a crystalline solid. A stock solution may be made by dissolving the dimethoxycurcumin in the solvent of choice, which should be purged with an inert gas. Dimethoxycurcumin is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of dimethoxycurcumin in these solvents is approximately 0.25, 5, and 10 mg/ml, respectively.

Dimethoxycurcumin is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, dimethoxycurcumin should first be dissolved in DMF and then diluted with the aqueous buffer of choice. Dimethoxycurcumin has a solubility of approximately 0.3 mg/ml in a 1:2 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Dimethoxycurcumin is a derivative of curcumin (Item No. 81025) that has anti-inflammatory and antioxidant activities.¹⁻³ It inhibits mitogen-induced proliferation of CD4⁺ T cells, CD8⁺ T cells, and B cells, as well as secretion of IL-2, IL-4, IL-6, and IFN-γ induced by concanavalin A (Item No. 14951) in isolated human lymphocytes when used at concentrations ranging from 1 to 10 μ M.¹ Dimethoxycurcumin (1-10 µM) inhibits LPS-induced nitric oxide (NO) production and expression of inducible nitric oxide synthase (iNOS) in RAW 264.7 cells.² It inhibits hemolysis of isolated human red blood cells (RBCs) induced by AAPH (Item No. 82235) when used at a concentration of 10 μ M.³

References

- 1. Patwardhan, R.S., Checker, R., Sharma, D., et al. Dimethoxycurcumin, a metabolically stable analogue of curcumin, exhibits anti-inflammatory activities in murine and human lymphocytes. Biochem. Pharmacol. 82(6), 642-657 (2011).
- 2. Pae, H.-O., Jeong, S.-O., Kim, H.S., et al. Dimethoxycurcumin, a synthetic curcumin analogue with higher metabolic stability, inhibits NO production, inducible NO synthase expression and NF-κB activation in RAW264.7 macrophages activated with LPS. Mol. Nutr. Food Res. 52(9), 1082-1091 (2008).
- 3. Deng, S.-L., Chen, W.-F., Zhou, B., et al. Protective effects of curcumin and its analogues against free radical-induced oxidative haemolysis of human red blood cells. Food Chem. 98(1), 112-119 (2006).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 07/02/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM