



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



(±)12(13)-DiHOME-d₄

Item No. 10009994

Formal Name: (±)12,13-dihydroxy-9Z-octadecenoic-9,10,12,13-d₄ acid
Synonym: Isoleukotoxin diol-d₄
MF: C₁₈H₃₀D₄O₄
FW: 318.5
Chemical Purity: ≥98% (±)12(13)-DiHOME
Deuterium Incorporation: ≥99% deuterated forms (d₁-d₄); ≤1% d₀
Supplied as: A solution in methyl acetate
Storage: -20°C
Stability: ≥1 year



NOTE: Relative stereochemistry shown in chemical structure

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

(±)12(13)-DiHOME-d₄ is intended for use as an internal standard for the quantification of (±)12(13)-DiHOME (Item No. 10009832) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

(±)12(13)-DiHOME-d₄ is supplied as a solution in methyl acetate. To change the solvent, simply evaporate the methyl acetate under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide purged with an inert gas can be used. The solubility of (±)12(13)-DiHOME-d₄ in these solvents is approximately 20 mg/ml.

Description

12(13)-DiHOME is the diol form of 12(13)-EpOME, a cytochrome P450-derived epoxide of linoleic acid also known as isoleukotoxin.¹ It is formed from 12(13)-EpOME by soluble epoxide hydrolase (sEH) in neutrophils.²

References

1. Greene, J.F., Williamson, K.C., Newman, J.W., et al. *Arch. Biochem. Biophys.* **376**(2), 420-432 (2000).
2. Moghaddam, M.F., Grant, D.F., Cheek, J.M., et al. *Nat. Med.* **3**(5), 562-566 (1997).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 04/05/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM