

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



AG-490

Item No. 10010311

CAS Registry No.: 133550-30-8

Formal Name: (2E)-2-cyano-3-(3,4-

dihydroxyphenyl)-N-(phenylmethyl)-

2-propenamide

Synonym: Tyrphostin AG-490

MF: $C_{17}H_{14}N_2O_3$ FW: 294.3 **Purity:** ≥98%

UV/Vis.: λ_{max} : 257, 363 nm Supplied as: A crystalline solid

-20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

HO

Laboratory Procedures

AG-490 is supplied as a crystalline solid. A stock solution may be made by dissolving the AG-490 in the solvent of choice, which should be purged with an inert gas. AG-490 is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of AG-490 in these solvents is approximately 10, 30, and 25 mg/ml, respectively.

AG-490 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, AG-490 should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. AG-490 has a solubility of approximately 0.1 mg/ml in a 1:10 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Protein tyrosine kinase (Ptk) inhibitors are potential antiproliferative agents for diseases caused by the hyperactivity of Ptks. Tyrphostins are a class of antiproliferative compounds which act as Ptk blockers.¹ Leukemic cells from patients in relapse have constitutively activated JAK2 Ptk. AG-490 is an inhibitor of JAK2 activity and selectively blocks leukemic cell growth in vitro and in vivo by inducing programmed cell death, with no deleterious effect on normal hematopoiesis. AG-490 almost completely blocks growth of all pre-B acute leukemia (ALL) cells at a concentration of 5 μM.² Furthermore, inhibition of JAK/STAT signaling in satellite cells via AG-490 and the STAT3 inhibitor 5,15-DPP (Item No. 16090) has been used to stimulate muscle regeneration in a model of aging skeletal muscle deterioration.³

References

- 1. Gazit, A., Yaish, P., Gilon, C., et al. Tyrphostins I: Synthesis and biological activity of protein tyrosine kinase inhibitors. J. Med. Chem. 32(10), 2344-2352 (1989).
- 2. Meydan, N., Grunberger, T., Dadi, H., et al. Inhibition of acute lymphoblastic leukaemia by a Jak-2 inhibitor. Nature 379(6566), 645-648 (1996).
- 3. Price, F.D., von Maltzahn, J., Bentzinger, C.F., et al. Inhibition of JAK-STAT signaling stimulates adult satellite cell function. Nat. Med. 20(10), 1174-1181 (2014).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 06/25/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM