

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



RbBP5 (human, recombinant)

Item No. 10947

Overview and Properties

RBQ3, Retinoblastoma binding Protein 5, SWD1, Set1c WD40 repeat protein, homolog Synonyms:

Source: Recombinant protein expressed in E. coli

Amino Acids: 2-538 (full length)

Uniprot No.: Q15291 Molecular Weight: 72.9 kDa

Storage: -80°C (as supplied)

Stability: ≥6 months

Purity: batch specific (≥80% estimated by SDS-PAGE)

50 mM Tris, pH 8.0, containing 150 mM sodium chloride and 20% glycerol Supplied in:


Endotoxin Testing: < 1.0 EU/μg, determined by the LAL endotoxin assay

Protein

batch specific mg/ml Concentration:

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Image



Lane 1: MW Markers Lane 2: RbBP5 (1 µg) Lane 3: RbBP5 (2 µg) Lane 4: RbBP5 (5 µg)

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/28/2024

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM

PRODUCT INFORMATION



Description

Retinoblastoma-binding protein 5 (RbBP5) is a ubiquitously expressed nuclear protein that contains WD40 repeat-like domains. RbBP5 binds directly to tumor suppressor retinoblastoma protein (RB) and regulates cell proliferation. Interaction of RbBP5 occurs preferentially with underphosphorylated RB through the E1A-binding pocket B.¹ RbBP5 is also an important component of the multi-subunit SET1 lysine methyltransferase protein complex, which includes MLL1.² RbBP5 interacts directly with WDR5 contributing to the activation of the MLL1 core protein complex.³-5 MLL1-4 protein complexes catalyze the di- and trimethylation of histone H3 at lysine 4 (H3K4me2/me3), leading to the maintenance of global H3K4 trimethylation.6

References

- 1. Saijo, M., Sakai, Y., Kishino, T., *et al.* Molecular cloning of a human protein that binds to the retinoblastoma protein and chromosomal mapping. *Genomics* **27**, 511-519 (1995).
- 2. Steward, M.M., Lee, J.S., O'Donovan, A., et al. Molecular regulation of H3K4 trimethylation by ASH2L, a shared subunit of MLL complexes. *Nat. Struct. Mol. Biol.* **13(9)**, 852-4 (2009).
- 3. Avdic, V., Zhang, P., Lanouette, S., et al. Structural and biochemical insights into MLL1 core complex assembly. Structure 19(1), 101-108 (2011).
- Odho, Z., Southall, S.M., and Wilson, J.R. Characterization of a novel WDR5-binding site that recruits RbBP5 through a conserved motif to enhance methylation of histone H3 lysine 4 by mixed lineage leukemia protein-1. J. Biol. Chem. 285(43), 32967-76 (2010).
- 5. Dou, Y., Milne, T.A., Ruthenburg, A.J., et al. Regulation of MLL1 H3K4 methyltransferase activity by its core components. *Nat. Struct. Mol. Biol.* **13(8)**, 713-9 (2006).
- Wang, P., Lin, C., Smith, E.R., et al. Global analysis of H3K4 methylation defines MLL family member targets and points to a role for MLL1-mediated H3K4 methylation in the regulation of transcriptional initiation by RNA polymerase II. Mol. Cell. Biol. 29(22), 6074-85 (2009).

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897