

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



SMI-4a

Item No. 11029

CAS Registry No.:	438190-29-5
Formal Name:	5Z-[[3-(trifluoromethyl)phenyl]
	methylene]-2,4-thiazolidinedione
MF:	C ₁₁ H ₆ F ₃ NO ₂ S
FW:	273.2 H—N _
Purity:	≥98%
UV/Vis.:	λ _{max} : 230, 320 nm Ó
Supplied as:	A crystalline solid CF ₃
Storage:	-20°C
Stability:	≥4 years
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.	

Laboratory Procedures

SMI-4a is supplied as a crystalline solid. A stock solution may be made by dissolving the SMI-4a in the solvent of choice, which should be purged with an inert gas. SMI-4a is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of SMI-4a in these solvents is approximately 30 mg/ml.

SMI-4a is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, SMI-4a should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. SMI-4a has a solubility of approximately 0.5 mg/ml in a 1:1 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

The serine/threonine Pim kinases have been suggested to promote the activity of the rapamycin-sensitive mammalian target of rapamycin (mTORC1), which regulates cell growth and survival.¹ Pim kinases are overexpressed in solid cancers and hematologic malignancies, and as such have become targets of small molecule inhibitors to prevent the progression of various cancers. SIM-4a is a Pim kinase inhibitor that blocks mTORC1 activity via activation of AMPK.¹ SIM-4a kills a wide range of both myeloid and lymphoid cell lines (with IC50 values ranging from 0.8 to 40 μ M).² Incubation of precursor T-cell lymphoblastic leukemia/lymphoma cells with 10 µM SMI-4a induces G₁ phase cell-cycle arrest, dose-dependent induction of p27Kip1, apoptosis through the mitochondrial pathway, and inhibition of the mTORC1 pathway.² In immunodeficient mice carrying subcutaneous pre-T-LBL tumors, treatment twice daily with 60 mg/kg SMI-4a causes a significant delay in the tumor growth.²

References

- 1. Beharry, Z., Mahajan, S., Zemskova, M., et al. The Pim protein kinases regulate energy metabolism and cell growth. Proc. Natl. Acad. Sci. USA 108(2), 528-533 (2011).
- 2. Lin, Y.W., Beharry, Z.M., Hill, E.G., et al. A small molecule inhibitor of Pim protein kinases blocks the growth of precursor T-cell lymphoblastic leukemia/lymphoma. Blood 115(4), 824-833 (2010).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/11/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM