



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



PRODUCT INFORMATION



GW 6471

Item No. 11697

CAS Registry No.: 880635-03-0

Formal Name: N-[(2S)-2-[[[(1Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propen-1-yl]amino]-3-[4-[2-(5-methyl-2-phenyl-4-oxazolyl)ethoxy]phenyl]propyl]-propanamide

MF: $C_{35}H_{36}F_3N_3O_4$


FW: 619.7

Purity: $\geq 98\%$

Stability: ≥ 2 years at -20°C

Supplied as: A crystalline solid

UV/Vis.: λ_{max} : 225, 278, 347, 352 nm



Laboratory Procedures

For long term storage, we suggest that GW 6471 be stored as supplied at -20°C . It should be stable for at least two years.

GW 6471 is supplied as a crystalline solid. A stock solution may be made by dissolving the GW 6471 in the solvent of choice. GW 6471 is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of GW 6471 in these solvents is approximately 10, 11, and 20 mg/ml, respectively.

GW 6471 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, GW 6471 should first be dissolved in DMF and then diluted with the aqueous buffer of choice. GW 6471 has a solubility of approximately 0.33 mg/ml in a 1:2 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

The peroxisome proliferator-activated receptor α (PPAR α) is a nuclear receptor that regulates the expression of genes involved in fatty acid metabolism, lipoprotein synthesis and metabolism, and inflammation.¹⁻³ GW 6471 is an antagonist of PPAR α (IC_{50} = 240 nM).⁴ It drives the displacement of coactivators from PPAR α and promotes the recruitment of co-repressor proteins like nuclear co-repressor.⁴ GW 6471 is used in cell-free, cell lysate, and whole cell systems to examine the impact of PPAR α antagonism.⁵⁻⁷

References

1. Kersten, S., Desvergne, B., and Wahli, W. *Nature* **405**, 421-424 (2000).
2. Tenenbaum, A., Motro, M., and Fisman, E.Z. *Cardiovasc. Diabetol.* **4**(14), 1-5 (2005).
3. Ahmed, W., Ziouzenkova, O., Brown, J., et al. *J. Intern. Med.* **262**, 184-198 (2007).
4. Xu, H.E., Stanley, T.B., Montana, V.G., et al. *Nature* **415**(6873), 813-817 (2002).
5. Wang, W., Lin, Q., Lin, R., et al. *Exp. Cell Res.* **319**(10), 1523-1533 (2013).
6. Downer, E.J., Clifford, E., Amu, S., et al. *J. Biol. Chem.* **287**(30), 25440-25453 (2012).
7. Müller, M.Q., de Koning, L.I., Schmidt, A., et al. *J. Med. Chem.* **52**(9), 2875-2879 (2009).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/23/2015

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM