

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Arteether

Item No. 11814

CAS Registry No.: 75887-54-6

Formal Name: (3R,5aS,6R,8aS,9R,10S,12R,12aR)-

> 10-ethoxydecahydro-3,6,9-trimethyl-3,12-epoxy-12H-pyrano[4,3-j]-1,2-

benzodioxepin

Synonyms: Artemotil, Dihydroartemisinin ethyl ether,


NSC 665971

MF: $C_{17}H_{28}O_5$ 312.4 FW: **Purity:** ≥95%

A crystalline solid Supplied as:

-20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Arteether is supplied as a crystalline solid. A stock solution may be made by dissolving the arteether in the solvent of choice. Arteether is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of arteether in these solvents is approximately 16, 10, and 20 mg/ml, respectively.

Arteether is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, arteether should first be dissolved in DMF and then diluted with the aqueous buffer of choice. Arteether has a solubility of approximately 0.5 mg/ml in a 1:1 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Arteether is a derivative of the antimalarial drug artemisinin (Item No. 11816). Arteether inhibits P. falciparum with an IC₅₀ value of 1.27 nM in vitro. It is effective against various strains of P. berghei in mice with ED_{90} values of 0.5-8.2 nM.

Reference

1. Brossi, A., Venugopalan, B., Dominguez Gerpe, L., et al. Arteether, a new antimalarial drug: Synthesis and antimalarial properties. J. Med. Chem. 31(3), 645-650 (1988).

WARNINGTHIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after ndling. Before use, the user <u>must</u> review the <u>complete</u> Safety Data Sheet, which has been sent *via* email to yo

WARRANTY AND LIMITATION OF REMEDY
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information

Copyright Cayman Chemical Company, 10/10/2017

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA **PHONE:** [800] 364-9897

[734] 971-3335 **FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM