

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Colchicoside

Item No. 11831

CAS Registry No.: 477-29-2

Formal Name: N-[(7S)-3-(β-D-glucopyranosyloxy)-

5,6,7,9-tetrahydro-1,2,10-trimethoxy-9-

oxobenzo[a]heptalen-7-yl]-acetamide

Synonym: 3-Demethylcolchicine glucoside


MF: $C_{27}H_{33}NO_{11}$ FW: 547.6 **Purity:** >98%

λ_{max}: 244, 247 nm UV/Vis.: Supplied as: A crystalline solid

-20°C Storage: Stability: ≥4 years

Item Origin: Natural/Source unknown

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Colchicoside is supplied as a crystalline solid. A stock solution may be made by dissolving the colchicoside in the solvent of choice, which should be purged with an inert gas. Colchicoside is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of colchicoside in these solvents is approximately 30 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of colchicoside can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of colchicoside in PBS, pH 7.2, is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Colchicoside is a glycoside form of 3-demethylcolchicine that has been found in G. superba. 1 Unlike cholchicine and 3-demethylcolchicine, colchicoside does not displace [3H]-colchicine from rat brain tubulin in vitro when used at a concentration of 25 μ M.² Colchicoside (1 and 10 μ M) increases levels of the cytochrome P450 (CYP) isoforms CYP2C9 and CYP2E1 in primary human hepatocytes.³

References

- 1. Capistrano, I.R., Vangestel, C., Vanpachtenbeke, H., et al. Coadministration of a Gloriosa superba extract improves the in vivo antitumoural activity of gemcitabine in a murine pancreatic tumour mode. Phytomedicine 23(12), 1434-1440 (2016).
- 2. Sugio, K., Maruyama, M., Tsurufuji, S., et al. Separation of tubulin-binding and anti-inflammatory activity in colchicine analogs and congeners. Life Sci. 40(1), 35-39 (1987).
- 3. Dvorák, Z., Ulrichová, J., Modrianský, M., et al. Effect of colchicine and its derivatives on the expression of selected isoforms of cytochrome P450 in primary cultures of human hepatocytes. Acta Univ. Palacki. Olomuc. Fac. Med. 143, 47-50 (2000).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 12/15/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM