



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



Ro 67-7476

Item No. 11993

CAS Registry No.: 298690-60-5
Formal Name: 2S-(4-fluorophenyl)-1-[(4-methylphenyl)sulfonyl]-pyrrolidine
MF: $C_{17}H_{18}FNO_2S$
FW: 319.4
Purity: $\geq 98\%$
UV/Vis.: λ_{max} : 231 nm
Supplied as: A crystalline solid
Storage: $-20^{\circ}C$
Stability: ≥ 2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Ro 67-7476 is supplied as a crystalline solid. A stock solution may be made by dissolving the Ro 67-7476 in the solvent of choice. Ro 67-7476 is soluble in the organic solvent DMSO, which should be purged with an inert gas, at a concentration of approximately 30 mg/ml.

Description

Ro 67-7476 is a positive allosteric modulator of metabotropic glutamate receptor 1 (mGluR1) that enhances glutamate-induced calcium release in HEK293 cells expressing rat mGluR1a ($EC_{50} = 1.74 \mu M$).¹ It is selective for rat mGluR1 over rat mGluR2, mGluR4, and mGluR8 at a concentration of 10 μM . Ro 67-7476 increases the amplitude of mGluR1 excitatory postsynaptic potentials (EPSCs) evoked by 2,3-dihydroxy-6-nitro-7-sulfamoylbenzoquinoline, picrotoxin, or AP5 in the Purkinje cells of rat cerebellar slices. It activates ERK1/2 phosphorylation in the absence of exogenously added glutamate ($EC_{50} = 163.3 nM$).²

References

1. Knoflach, F., Mutel, V., Jolidon, S., *et al.* Positive allosteric modulators of metabotropic glutamate 1 receptor: Characterization, mechanism of action, and binding site. *Proc. Natl. Acad. Sci. U.S.A.* **98(23)**, 13402-13407 (2001).
2. Sheffler, D.J. and Conn, P.J. Allosteric potentiators of metabotropic glutamate receptor subtype 1a differentially modulate independent signaling pathways in baby hamster kidney cells. *Neuropharmacology* **55(4)**, 419-427 (2008).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/22/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM