

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



TCN 238

Item No. 11994

CAS Registry No.:	125404-04-8	
Formal Name:	(E)-4-(2-phenylethenyl)-2-pyrimidinamine	
MF:	C ₁₂ H ₁₁ N ₃	N II
FW:	197.2	
Purity:	≥98%	N NH ₂
UV/Vis.:	λ _{max} : 228, 295 nm	
Supplied as:	A crystalline solid	
Storage:	-20°C	~
Stability:	≥2 years	

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

TCN 238 is supplied as a crystalline solid. A stock solution may be made by dissolving the TCN 238 in the solvent of choice. TCN 238 is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of TCN 238 in ethanol is approximately 5 mg/ml and approximately 25 mg/ml in DMSO and DMF.

TCN 238 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, TCN 238 should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. TCN 238 has a solubility of approximately 0.5 mg/ml in a 1:1 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

TCN 238 is an orally bioavailable positive allosteric modulator of the metabotropic glutamate receptor 4 (mGluR4; EC₅₀ = 1 μ M).¹ It has a greater than 30-fold selectivity for mGluR4 over mGluR5 using human recombinant receptors in CHO-K1 cells. It has no activity at the adenosine A2A, serotonin 5-HT1A, or a_{2A} -adrenergic receptors at concentrations up to 10 μ M. In a haloperidol-induced catalepsy rat model of Parkinson's disease, it dose-dependently reduces catalepsy with an ED_{50} of approximately 1 mg/kg. TCN 238 administration in rats leads to downregulation of the mGluR4 gene, GRM4, in the hippocampus and the gene for the GABA_{Δ} receptor α subunit, GABRA1, in the frontal cortex without affecting hippocampal-dependent memory.²

References

- 1. East, S.P., Bamford, S.J., Dietz, M.G.A., et al. An orally bioavailable positive allosteric modulator of the mGlu4 receptor with efficacy in an animal model of motor dysfunction. Bioorg. Med. Chem. Lett. 20(16), 4901-4905 (2010).
- 2. Pershina, E.V., and Arkhipov, V.I. Subacute activation of mGlu4 receptors causes the feedback inhibition of its gene expression in rat brain. Life Sci. 153, 50-54 (2016).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 08/29/2017

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM