

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



BI-6015

Item No. 12032

CAS Registry No.:	93987-29-2	
Formal Name:	2-methyl-1-[(2-methyl-5-nitrophenyl)	
	sulfonyl]-1H-benzimidazole	
MF:	$C_{15}H_{13}N_3O_4S$	
FW:	331.3	
Purity:	≥95% 0=s	
UV/Vis.:	λ_{max} : 245 nm	
Supplied as:	A crystalline solid	
Storage:	-20°C	
Stability:	≥4 years	
Information represent	s the product specifications. Batch specific analytical results are provided on each certificate of analys	is.

Laboratory Procedures

BI-6015 is supplied as a crystalline solid. A stock solution may be made by dissolving the BI-6015 in the solvent of choice, which should be purged with an inert gas. BI-6015 is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of BI-6015 in these solvents is approximately 14 and 16 mg/ml, respectively.

BI-6015 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, BI-6015 should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. BI-6015 has a solubility of approximately 0.09 mg/ml in a 1:10 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Hepatocyte nuclear factor 4α (HNF4 α) is a nuclear receptor transcription factor that controls the expression of many genes including those involved in glucose and lipid homeostasis and maintenance of epithelial differentiation. BI-6015 is a small molecule antagonist of HNF4 α that has been shown to repress the expression of known HNF4 α target genes.¹ At concentrations up to 20 μ M, BI-6015 has been shown to reduce endogenous insulin gene expression by as much as 50-fold in T6PNE cells.¹ The hepatocytes from livers of mice injected with 10 and 30 mg/kg BI-6015 exhibit marked, dose-dependent fat accumulation (steatosis).¹ Additionally, at a daily dose of 30 mg/kg, BI-6015 induces apoptosis in a human hepatocellular carcinoma mouse model.¹

Reference

1. Kiselyuk, A., Lee, S.-H., Farber-Katz, S., et al. ΗΝF4α antagonists discovered by a high-throughput screen for modulators of the human insulin promoter. Chem. Biol. 19(7), 806-818 (2012).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/26/2024

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM