



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



Gefitinib

Item No. 13166

CAS Registry No.: 184475-35-2
Formal Name: N-(3-chloro-4-fluorophenyl)-7-methoxy-6-[3-(4-morpholinyl)propoxy]-4-quinazolinamine
Synonym: ZD 1839
MF: C₂₂H₂₄ClFN₄O₃
FW: 446.9
Purity: ≥98%
UV/Vis.: λ_{max}: 205, 226, 250, 332 nm
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Gefitinib is supplied as a crystalline solid. A stock solution may be made by dissolving the gefitinib in the solvent of choice. Gefitinib is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of gefitinib in ethanol is approximately 0.3 mg/ml and approximately 20 mg/ml in DMSO and DMF.

Gefitinib is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, gefitinib should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. Gefitinib has a solubility of approximately 0.5 mg/ml in a 1:1 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Gefitinib is an EGFR inhibitor (IC₅₀s = 0.023-0.079 μM).² It inhibits colony formation of GEO colon, ZR-75-1 and MCF-10A breast, and OVCAR-3 ovarian cancer cell lines in soft agar assays (IC₅₀s = 0.2-0.4 μM).¹ Gefitinib (0.1 to 1 μM) induces apoptosis and inhibits EGFR autophosphorylation in the same cells. *In vivo*, gefitinib (1.25, 2.5, and 5 mg/kg) reduces tumor volume and increases survival in a GEO mouse xenograft model. Formulations containing gefitinib have been used in the treatment of non-small cell lung cancer (NSCLC).

References

1. Ciardiello, F., Caputo, R., Bianco, R., *et al.* Antitumor effect and potentiation of cytotoxic drugs activity in human cancer cells by ZD-1839 (Iressa), an epidermal growth factor receptor-selective tyrosine kinase inhibitor. *Clin. Cancer Res.* **6**(5), 2053-2063 (2000).
2. Mendelsohn, J. and Baselga, J. The EGF receptor family as targets for cancer therapy. *Oncogene* **19**(56), 6550-6565 (2000).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/14/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM