



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION

G-418 (sulfate)

Item No. 13200

CAS Registry No.: 108321-42-2

Formal Name: O-2-amino-2,7-dideoxy-D-glycero- α -D-glucopyranosyl-(1 $\rightarrow$ 4)-O-[3-deoxy-4-C-methyl-3-(methylamino)- β -L-arabinopyranosyl-(1 $\rightarrow$ 6)]-2-deoxy-D-streptamine, disulfate

MF: $C_{20}H_{40}N_4O_{10} \cdot 2H_2SO_4$


FW: 692.7

Purity: $\geq 95\%$

Supplied as: A crystalline solid

Storage: $-20^\circ C$

Stability: ≥ 4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

G-418 (sulfate) is supplied as a crystalline solid. Aqueous solutions of G-418 (sulfate) can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of G-418 (sulfate) in PBS (pH 7.2) is approximately 100 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

G-418 is an aminoglycoside antibiotic.¹ It is active against *T. thermophila* ($IC_{50} = 10 \mu M$). It inhibits the 80S ribosome and protein elongation during translation in a cell-free assay when used at a concentration of 2 nM.² G-418 is used in molecular biology as a selective agent in cell lines or plasmids containing the neomycin resistance gene, *neoR*.³ It increases survival in mouse models of *Salmonella*, *P. vulgaris*, *K. pneumoniae*, or *E. coli* when administered at doses ranging from 0.8 to 4 mg/kg.⁴

References

1. Eustice, D.C. and Wilhelm, J.M. Mechanisms of action of aminoglycoside antibiotics in eucaryotic protein synthesis. *Antimicrob. Agents Chemother.* **26**(1), 53-60 (1984).
2. Bar-Nun, S., Shneyour, Y., and Beckmann, J.S. G-418, an elongation inhibitor of 80 S ribosomes. *Biochim. Biophys. Acta* **741**(1), 123-127 (1983).
3. Giordano-Santini, R., Milstein, S., Svrzikapa, N., et al. An antibiotic selection marker for nematode transgenesis. *Nat. Methods* **7**(9), 721-723 (2010).
4. Waitz, J.A., Sabatelli, F., Menzel, F., et al. Biological activity of antibiotic G-418, a new *Micromonospora*-produced aminoglycoside with activity against protozoa and helminths. *Antimicrob. Agents Chemother.* **6**(5), 579-581 (1974).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/07/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM