



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



IWP-4

Item No. 13954

CAS Registry No.: 686772-17-8

Formal Name: N-(6-methyl-2-benzothiazolyl)-
2-[[3,4,6,7-tetrahydro-
3-(2-methoxyphenyl)-4-
oxothieno[3,2-d]pyrimidin-2-yl]
thio]-acetamide

Synonym: Inhibitor of Wnt Production-4

MF: $C_{23}H_{20}N_4O_3S_3$

FW: 496.6


Purity: $\geq 95\%$

UV/Vis.: λ_{max} : 279, 302, 343 nm

Supplied as: A crystalline solid

Storage: -20°C

Stability: ≥ 2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

IWP-4 is supplied as a crystalline solid. A stock solution may be made by dissolving the IWP-4 in the solvent of choice, which should be purged with an inert gas. IWP-4 is soluble in organic solvents such as DMSO and dimethyl formamide (DMF). The solubility of IWP-4 in these solvents is approximately 2 and 5 mg/ml, respectively.

IWP-4 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, IWP-4 should first be dissolved in DMF and then diluted with the aqueous buffer of choice. IWP-4 has a solubility of approximately 0.3 mg/ml in a 1:2 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

IWP-4 is an inhibitor of Wnt production that impairs Wnt pathway activity *in vitro* with an IC_{50} value of 25 nM.¹ IWP-4 inactivates Porcupine, a membrane-bound O-acyltransferase responsible for palmitoylating Wnt proteins, which is essential for their signaling ability and secretion.¹ At 5 μM , IWP-4 has been shown to block Wnt-dependent phosphorylation of the low-density lipoprotein receptor-related protein 6 receptor and the scaffold protein Dishevelled, preventing the accumulation of β -catenin.¹ This compound has been used to induce cardiomyocyte differentiation from human pluripotent stem cells.²

References

1. Chen, B., Dodge, M.E., Tang, W., *et al.* Small molecule-mediated disruption of Wnt-dependent signaling in tissue regeneration and cancer. *Nat. Chem. Biol.* **5**(2), 100-107 (2009).
2. Lian, X., Hsiao, C., Wilson, G., *et al.* Robust cardiomyocyte differentiation from human pluripotent stem cells via temporal modulation of canonical Wnt signaling. *Proc. Natl. Acad. Sci. USA* E1848-E1857 (2012).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 06/26/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM