



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



# PRODUCT INFORMATION



## Prostaglandin E<sub>2</sub>-d<sub>4</sub>-1-glycerol ester

Item No. 14029

**Formal Name:** 9-oxo-11 $\alpha$ ,15S-dihydroxy-prosta-5Z,13E-dien-1-oic-2,2,3,3-d<sub>4</sub> acid, 1-glycerol ester

**Synonym:** PGE<sub>2</sub>-d<sub>4</sub>-1-glycerol ester

**MF:** C<sub>23</sub>H<sub>34</sub>D<sub>4</sub>O<sub>7</sub>

**FW:** 430.6

**Chemical Purity:** ≥98% (isomeric mixture)  
(Prostaglandin E<sub>2</sub>-1-glycerol ester)


**Deuterium**

**Incorporation:** ≥99% deuterated forms (d<sub>1</sub>-d<sub>4</sub>); ≤1% d<sub>0</sub>

**Supplied as:** A solution in acetonitrile

**Storage:** -20°C

**Stability:** ≥1 year



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

Prostaglandin E<sub>2</sub>-d<sub>4</sub>-1-glycerol ester (PGE<sub>2</sub>-d<sub>4</sub>-1-glycerol ester) is intended for use as an internal standard for the quantification of PGE<sub>2</sub>-1-glycerol ester (Item No. 10140) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

PGE<sub>2</sub>-d<sub>4</sub>-1-glycerol ester is supplied as a solution in acetonitrile. To change the solvent, simply evaporate the acetonitrile under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide purged with an inert gas can be used. The solubility of PGE<sub>2</sub>-d<sub>4</sub>-1-glycerol ester in these solvents is approximately 10 mg/ml.

### Description

2-Arachidonoyl glycerol (2-AG) has been isolated from porcine brain, and has been characterized as the natural endocannabinoid ligand for the CB<sub>1</sub> receptor.<sup>1,2</sup> Incubation of 2-AG with COX-2 and specific prostaglandin H<sub>2</sub> (PGH<sub>2</sub>) isomerases in cell cultures and isolated enzyme preparations results in prostaglandin glycerol ester formation.<sup>3</sup> The biosynthesis of PGH, PGD, PGE, PGF, and TXA-2-glycerol ester compounds have all been documented. The 2-glycerol ester moiety equilibrates rapidly (within minutes) with the more stable 1-glycerol ester, producing a 10:90 2:1-glycerol ester mixture in typical aqueous media. While the stability and metabolism of these prostaglandin products has been investigated, little is known about their intrinsic biological activity.<sup>4</sup>

### References

1. Sugiura, T., Kodaka, T., Kondo, S., *et al.* 2-Arachidonoylglycerol, a putative endogenous cannabinoid receptor ligand, induces rapid, transient elevation of intracellular free Ca<sup>2+</sup> in neuroblastoma X glioma hybrid NG108-15 cells. *Biochem. Biophys. Res. Commun.* **229**(1), 58-64 (1996).
2. Sugiura, T., Kodaka, T., Kondo, S., *et al.* Is the cannabinoid CB<sub>1</sub> receptor a 2-arachidonoylglycerol receptor? Structural requirements for triggering a Ca<sup>2+</sup> transient in NG108-15 cells. *J. Biochem.* **122**(4), 890-895 (1997).
3. Kozak, K.R., Crews, B.C., Morrow, J.D., *et al.* Metabolism of the endocannabinoids, 2-arachidonoylglycerol and anandamide, into prostaglandin, thromboxane, and prostacyclin glycerol esters and ethanolamides. *J. Biol. Chem.* **277**(47), 44877-44885 (2002).
4. Kozak, K.R., Crews, B.C., Ray, J.L., *et al.* Metabolism of prostaglandin glycerol esters and prostaglandin ethanolamides *in vitro* and *in vivo*. *J. Biol. Chem.* **276**(40), 36993-36998 (2001).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 03/30/2023

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM