

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



S18886

Item No. 14059

CAS Registry No.: 165538-40-9

Formal Name: (6R)-6-[[(4-chlorophenyl)sulfonyl]


amino]-5,6,7,8-tetrahydro-2-methyl-

1-naphthalenepropanoic

Synonyms: Terutroban, Triplion C₂₀H₂₂CINO₄S MF:

FW: 407.9 Purity: ≥98%

 λ_{max} : 223 nm UV/Vis.: Supplied as: A crystalline solid



Laboratory Procedures

S18886 is supplied as a crystalline solid. A stock solution may be made by dissolving the S18886 in the solvent of choice. S18886 is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of \$18886 in ethanol is approximately 2 mg/ml, and approximately 30 mg/ml in DMSO and DMF.

S18886 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, S18886 should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. S18886 has a solubility of approximately 0.5 mg/ml in a 1:1 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

The lipid mediator thromboxane A2 (TXA2) plays a key role in platelet aggregation and vascular and bronchial smooth muscle constriction. The actions of TXA2 are mediated by its specific G protein coupled-receptor, referred to as the TXA_2 receptor or TP. S18886 is an orally active, long-acting antagonist of the TP receptor with an IC₅₀ value of 16.4 nM.¹ It inhibits TP receptor-mediated vascular contractions, platelet aggregation, and adhesion and infiltration of monocytes/macrophages in various pre-clinical and clinical models of thrombosis or atherosclerosis.²⁻⁵ S18886 is the active isomer of the potent TP receptor antagonist, S18204.6 Antagonists of TP receptors have advantages over aspirin (Item No. 70260) as they not only block the effect of TXA2 on platelets, but also inhibit other ligands such as prostaglandin endoperoxides and isoprostanes.^{2,5}

References

- 1. Wang, X., Liu, Li., Huang, L., et al. ACS Med. Chem. Lett. 5(9), 1015-1020 (2014).
- 2. Jones, R.L., Giembycz, M.A., and Woodward, D.F. Br. J. Pharmacol. 158(1), 104-145 (2009).
- 3. Hong, T.-T., Huang, J., Driscoll, E., et al. J. Cardiovasc. Pharmacol. 48(5), 239-248 (2006).
- 4. Osende, J.I., Shimbo, D., Fuster, V., et al. J. Thromb. Haemost. 2(3), 492-498 (2004).
- 5. Cayatte, A.J., Du, Y., Oliver-Krasinski, J., et al. Artioscler. Thromb. Vasc. Biol. 20(7), 1724-1728 (2000).
- Simonet, S., Descombes, J.-J., Vallez, M.O., et al. Adv. Exp. Med. Biol. 433, 173-176 (1997).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

al should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 01/17/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM