

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



L-Mimosine

Item No. 14337

CAS Registry No.: 500-44-7

Formal Name: (αS)-amino-3-hydroxy-4-oxo-

1(4H)-pyridinepropanoic acid


Synonyms: Leucenol, NSC 69188

MF: $C_8H_{10}N_2O_4$ 198.2 FW: ≥98% **Purity:**

 λ_{max} : 216, 283 nm UV/Vis.: Supplied as: A crystalline solid

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

L-Mimosine is supplied as a crystalline solid. L-Mimosine is sparingly soluble in aqueous solutions. To enhance aqueous solubility, dilute the organic solvent solution into aqueous buffers or isotonic saline. If performing biological experiments, ensure the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. We do not recommend storing the aqueous solution for more than one day.

Description

L-Mimosine is a non-protein amino acid that can be isolated from certain plants and fungi. It chelates iron and copper and has been shown to reduce iron overload in animal models.^{1,2} L-Mimosine inhibits certain enzymes that contain iron or copper, including arginase (IC $_{50}$ = 3.7 μ M), polyphenoloxidase, and dopamine hydroxylase.²⁻⁴ It also inhibits the iron-containing enzyme deoxyhypusine hydroxlase, preventing the synthesis of the eukaryotic initiation factor 5A and blocking cell cycling.⁵

References

- 1. Kontoghiorghe, C. N., Kolnagou, A., and Kontoghiorghes, G. J. Phytochelators intended for clinical use in iron overload, other diseases of iron imbalance and free radical pathway. Molecules 20, 20841-20872 (2015).
- 2. Nirmal, N. P., Benjakul, S. Inhibitory effects of mimosine on polyphenoloxidase from cephalothoraxes of Pacific white shrimp (Litopenaeus vannamei). J. Agric. Food Chem. 59(18), 10256-10260 (2011).
- Ahmad, V. U., Ullah, F., Hussain, J., et al. Tyrosinase inhibitors from Rhododendron collettianum and their structure - activity relationship (SAR) studies. Chem. Pharm. Bull. 52(12), 1458-1461 (2004).
- Hashiguchi, H. and Takahashi, H. Inhibition of two copper-containing enzymes, tyrosinase and dopamine β-hydroxylase, by L-mimosine. Mol. Pharmacol 13(2), 362-367 (1977).
- Hanauske-Abel, H. M., Park, M. H., Hanauske, A. R., et al. Inhibition of the G1-S transition of the cell cycle by inhibitors of deoxyhypusine hydroxylation. Biochim. Biophys. Acta. 1221(2), 115-124 (1994).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

al should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/07/2017

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM