

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Gly-Phe-\u03b3-naphthylamide

Item No. 14634

CAS Registry No.: 21438-66-4

glycyl-N-2-naphthalenyl-L-Formal Name:

phenylalaninamide


Synonym: H-Gly-Phe-BNA MF: $C_{21}H_{21}N_3O_2$ 347.4 FW:

≥95% **Purity:**

UV/Vis.: λ_{max} : 243, 283 nm Supplied as: A crystalline solid

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Gly-Phe-β-naphthylamide is supplied as a crystalline solid. A stock solution may be made by dissolving the Gly-Phe-\u00d3-naphthylamide in the solvent of choice, which should be purged with an inert gas. Gly-Phe-β-naphthylamide is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of Gly-Phe-β-naphthylamide in these solvents is approximately 1, 20, and 25 mg/ml, respectively.

Gly-Phe-β-naphthylamide is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, Gly-Phe-β-naphthylamide should first be dissolved in DMF and then diluted with the aqueous buffer of choice. Gly-Phe-β-naphthylamide has a solubility of approximately 0.5 mg/ml in a 1:1 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Gly-Phe-β-naphthylamide is a substrate of cathepsin C that accumulates within the lysosome. Hydrolysis by cathepsin C degrades Gly-Phe-β-naphthylamide into fragments that do not easily diffuse through the lysosomal membrane, leading to a loss of lysosome membrane integrity. This compound has been used to study intralysosomal hydrolysis, lysosomal membrane permeability, and the function of cathepsin C.^{1,2} At 50 μM, Gly-Phe- β -naphthylamide can inhibit the cathepsin-dependent activation of caspase-8.³

References

- 1. Jadot, M., Colmant, C., Wattiaux-De Coninck, S., et al. Intralysosomal hydrolysis of glycyl-L-phenylalanine 2-naphthylamide. Biochem. J. 219(3), 965-970 (1984).
- 2. Qiu, G.F., Feng, H.Y., and Yamano, K. Expression and purification of active recombinant cathepsin C (dipeptidyl aminopeptidase I) of kuruma prawn Marsupenaeus japonicus in insect cells. J. Biomed. Biotechnol. 746289 (2009).
- 3. Baumgartner, H.K., Gerasimenko, J.V., Thorne, C., et al. Caspase-8-mediated apoptosis induced by oxidative stress is independent of the intrinsic pathway and dependent on cathepsins. Am. J. Physiol. Gastrointest. Liver Physiol. 293(1), 296-307 (2007).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/17/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM