



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# PRODUCT INFORMATION



## Quinacrine (hydrochloride hydrate)

Item No. 15041

**Formal Name:** N<sup>4</sup>-(6-chloro-2-methoxyacridin-9-yl)-N<sup>1</sup>,N<sup>1</sup>-diethylpentane-1,4-diamine, dihydrochloride hydrate

**Synonyms:** Atebrine, Mepacrine

**MF:** C<sub>23</sub>H<sub>30</sub>ClN<sub>3</sub>O • 2HCl [XH<sub>2</sub>O]

**FW:** 472.9

**Purity:** ≥95%


**UV/Vis.:** λ<sub>max</sub>: 223, 283, 346, 427, 449 nm

**Ex./Em. Max:** 436/525 nm (DNA and RNA);  
458/470-580 nm (acidic vesicles/lysosomes)

**Supplied as:** A crystalline solid

**Storage:** -20°C

**Stability:** ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

Quinacrine (hydrochloride hydrate) is supplied as a crystalline solid. A stock solution may be made by dissolving the quinacrine (hydrochloride hydrate) in the solvent of choice. Quinacrine (hydrochloride hydrate) is soluble in organic solvents such as ethanol and DMSO, which should be purged with an inert gas. The solubility of quinacrine (hydrochloride hydrate) in these solvents is approximately 30 and 5 mg/ml, respectively.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of quinacrine (hydrochloride hydrate) can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of quinacrine (hydrochloride hydrate) in PBS, pH 7.2, is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

### Description

Quinacrine is an acridine compound with diverse biological activities that has also been used as a stain for DNA and RNA in fixed cells and for fluorescent tracking of acidic vesicles, such as lysosomes, in live cells.<sup>1-4</sup> It has anti-protozoal properties, potently prevents misfolding of prion protein (EC<sub>50</sub> = 0.3 μM), blocks voltage-dependent sodium channels (IC<sub>50</sub> = 3.3 μM), and inhibits aldehyde oxidase (IC<sub>50</sub> = 3.3 μM).<sup>1-4</sup> Quinacrine is also an effective riboflavin antagonist, associating with the riboflavin-binding protein (K<sub>i</sub> = 6.7 μM), and an inhibitor of P-glycoprotein (EC<sub>50</sub> = 14.4 μM).<sup>5,6</sup> At much higher doses, quinacrine inhibits phospholipase A<sub>2</sub> activity.<sup>7,8</sup> Quinacrine binds to nucleic acids and has been used in fixed cells to stain DNA and RNA with excitation/emission maxima of 436/525 nm, respectively.<sup>9</sup> In live cells, it is taken up into acidic vesicles, such as lysosomes, and can be used to for long-term imaging with an excitation of 458 nm and emission in the 470-580 nm range.<sup>10</sup>

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 03/20/2018

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897

[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
[WWW.CAYMANCHEM.COM](http://WWW.CAYMANCHEM.COM)

# PRODUCT INFORMATION



## References

1. Upcroft, P. and Upcroft, J.A. Drug targets and mechanisms of resistance in the anaerobic protozoa. *Clin. Microbiol. Rev.* **14**(1), 150-164 (2001).
2. Vogtherr, M., Grimme, S., Elshorst, B., *et al.* Antimalarial drug quinacrine binds to C-terminal helix of cellular prion protein. *J. Med. Chem.* **46**(17), 3563-3564 (2003).
3. McNeal, E.T., Lewandowski, G.A., Daly, J.W., *et al.* [<sup>3</sup>H]Batrachotoxinin A 20 $\alpha$ -benzoate binding to voltage-sensitive sodium channels: A rapid and quantitative assay for local anesthetic activity in a variety of drugs. *J. Med. Chem.* **28**(3), 381-388 (1985).
4. Pryde, D.C., Dalvie, D., Hu, Q., *et al.* Aldehyde oxidase: An enzyme of emerging importance in drug discovery. *J. Med. Chem.* **53**(24), 8441-8460 (2010).
5. Plantinga, A., Witte, A., Li, M.H., *et al.* Bioanalytical screening of riboflavin antagonists for targeted drug delivery - A thermodynamic and kinetic study. *ACS Med. Chem. Lett.* **2**(5), 363-367 (2011).
6. Tiberghien, F. and Loor, F. Ranking of P-glycoprotein substrates and inhibitors by a calcein-AM fluorometry screening assay. *Anti-Cancer Drugs* **7**(5), 568-578 (1996).
7. Farooqui, A.A., Ong, W.Y., and Horrocks, L.A. Inhibitors of brain phospholipase A<sub>2</sub> activity: Their neuropharmacological effects and therapeutic importance for the treatment of neurologic disorders. *Pharmacol. Rev.* **58**(3), 591-620 (2006).
8. Caro, A.A. and Cederbaum, A.I. Role of phospholipase A<sub>2</sub> activation and calcium in CYP2E1-dependent toxicity in HepG2 cells. *J. Biol. Chem.* **278**(36), 33866-33877 (2003).
9. Kasten, F.H. Table 2.3. *Fluorescent and Luminescent Probes for Biological Activity: A Practical Guide to Technology for Quantitative Real-Time Analysis* 33 (1999).
10. Pierzyńska-Mach, A., Janowski, P.A., and Dobrucki, J.W. Evaluation of acridine orange, LysoTracker Red, and quinacrine as fluorescent probes for long-term tracking of acidic vesicles. *Cytometry A.* **85**(8), 729-737 (2014).

## CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM