

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION **BQCA**



Item No. 15393

CAS Registry No.:	338747-41-4
Formal Name:	1,4-dihydro-1-[(4- o
	methoxyphenyl)methyl]-4-oxo-3- quinolinecarboxylic acid
MF:	C ₁₈ H ₁₅ NO ₄
FW:	309.3
Purity:	≥98%
UV/Vis.:	λ _{max} : 224, 315, 328 nm
Supplied as:	A crystalline solid
Storage:	-20°C
Stability:	As supplied, 2 years from the QC date provided on the Certificate of Analysis, when stored properly

Laboratory Procedures

BQCA is supplied as a crystalline solid. A stock solution may be made by dissolving the BQCA in the solvent of choice. BQCA is soluble in organic solvents such as DMSO and dimethyl formamide, which should be purged with an inert gas. The solubility of BQCA in these solvents is approximately 0.2 mg/ml.

BQCA is sparingly soluble in aqueous solutions. To enhance aqueous solubility, dilute the organic solvent solution into aqueous buffers or isotonic saline. If performing biological experiments, ensure the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. We do not recommend storing the aqueous solution for more than one day.

Description

BQCA is a highly selective positive allosteric modulator of the M_1 muscarinic acetylcholine receptor (mAChR), as it dose-dependently reduces the concentration of acetylcholine required to activate the M1 receptor.¹ The effective range for potentiation of M_1 in cells by BQCA is 0.1 to 100 μ M with an inflection point value of 845 nM when 3 nM acetylcholine is used.¹ BQCA displays no potentiation, agonism, or antagonism at other mAChRs at concentrations up to 100 μ M.¹ It has excellent brain penetration and increases the firing rate of medial prefrontal cortex neurons in vivo in rats.² BQCA prevents scopolamineinduced memory deficits in both contextual fear conditioning and a spontaneous alternation task in mice.^{1,3} It also restores impairment in reversal learning in a mouse model of Alzheimer's disease and improves memory performance in rats.^{2,4}

References

- 1. Ma, L., Seager, M.A., Wittman, M., et al. Proc. Natl. Acad. Sci. USA 106(37), 15950-15955 (2009).
- 2. Shirey, J.K., Brady, A.E., Jones, P.J., et al. J. Neurosci. 29(45), 14271-14286 (2009).
- 3. Chambon, C., Jatzke, C., Wegener, N., et al. Eur. J. Pharmacol. 697(1-3), 73-80 (2012).
- 4. Galloway, C.R., Lebois, E.P., Shagarabi, S., et al. Pharmacology 93(1-2), 57-64 (2014).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

al should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 04/27/2016

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA **PHONE:** [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM