

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Miconazole (nitrate)

Item No. 15420

CAS Registry No.: Formal Name:	22832-87-7 1-[2-(2,4-dichlorophenyl)-2-[(2,4- dichlorophenyl)methoxy]ethyl]- 1H-imidazole mononitrate	CI
Synonyms: MF: FW: Purity: Supplied as: Storage: Stability:	NSC 169434, R 14889 $C_{18}H_{14}Cl_4N_2O \bullet HNO_3$ 479.1 ≥98% A crystalline solid -20°C ≥2 years	

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Miconazole (nitrate) is supplied as a crystalline solid. A stock solution may be made by dissolving the miconazole (nitrate) in the solvent of choice. Miconazole (nitrate) is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of miconazole (nitrate) in ethanol is approximately 0.1 mg/ml and approximately 25 mg/ml in DMSO and DMF.

Miconazole (nitrate) is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, miconazole (nitrate) should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. Miconazole (nitrate) has a solubility of approximately 0.3 mg/ml in a 1:2 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Miconazole is a widely used antifungal imidazole that acts by inhibiting the 14α -demethylation of lanosterol, which consequently leads to the inhibition of ergosterol synthesis in fungal cell membranes.^{1,2} Additionally, miconazole has been shown to induce reactive oxygen species in fungal biofilms, demonstrating a MIC value of $\geq 256 \,\mu\text{M}$ against Candida spp.³

References

- 1. Vanden Bossche, H., Marichal, P., Gorrens, J., et al. Biochemical approaches to selective antifungal activity. Focus on azole antifungals. Mycoses 32(Suppl 1), 35-52 (1989).
- 2. Saag, M.S. and Dismukes, W.E. Azole antifungal agents: Emphasis on new triazoles. Antimicrob. Agents Chemother. 32(1), 1-8 (1988).
- 3. Delattin, N., Cammue, B.P.A., and Thevissen, K. Reactive oxygen species-inducing antifungal agents and their activity against fungal biofilms. Future Med. Chem. 6(1), 77-90 (2014).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/08/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM