

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



bpV(HOpic) (potassium salt) (technical grade)

Item No. 15438

CAS Registry No.: 722494-26-0

(5-hydroxy-2-pyridinecarboxylato- κN^1 , κO^2) Formal Name:

oxodiperoxy-vanadate(2-), dipotassium

Synonym: Bisperoxovanadium(HOpic)

MF: $C_6H_4NO_8V \bullet 2K$

FW: 347.2 **Purity:** ≥80%

Supplied as: A crystalline solid

Storage: -20°C Stability: ≥4 vears

Special Conditions: Product not stable in water. Decomposition starts immediately after being dissolved.

Rate of decomposition is slower in dry DMSO.

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

bpV(HOpic) (potassium salt) (technical grade) is supplied as a crystalline solid. A stock solution may be made by dissolving the bpV(HOpic) (potassium salt) (technical grade) in the solvent of choice, which should be purged with an inert gas. bpV(HOpic) (potassium salt) (technical grade) is soluble in organic solvents such as DMSO. It is also soluble in water. The solubility of bpV(HOpic) (potassium salt) (technical grade) in DMSO and water is approximately 4 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

bpV(HOpic) is a bisperoxovanadium (bpV) compound that inhibits several different protein tyrosine phosphatases (PTPs), with selectivity for PTEN (IC $_{50}$ = 14 nM). It also inhibits the vascular endothelial PTP, PTP- β (IC $_{50}$ = 4.9 μ M), and PTP-1 β B (IC $_{50}$ = 25.3 μ M). At 15 μ M, bpV(HOpic) has been shown to protect against stimulated ischemia-reperfusion injury in vitro by decreasing apoptosis and improving cell viability through the upregulation of the PI3K/Akt/eNOS/ERK prosurvival pathway.

References

- 1. Schmid, A.C., Byrne, R.D., Vilar, R., et al. Bisperoxovanadium compounds are potent PTEN inhibitors. FEBS Lett. 566(1-3), 35-38 (2004).
- 2. Keyes, K.T., Xu, J., Long, B., et al. Pharmacological inhibition of PTEN limits myocardial infarct size and improves left ventricular function postinfarction. Am. J. Physiol. Heart Circ. Physiol. 298(4), H1198-H1208 (2010).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 02/04/2025

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM