



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



COX-1 (mouse) Polyclonal Antibody

Item No. 160109

Overview and Properties

Contents:	This vial contains 500 µl of peptide affinity-purified polyclonal antibody.
Synonyms:	Cyclooxygenase 1, PGHS-1, Prostaglandin Endoperoxide Synthase 1, Prostaglandin G/H Synthase 1, Prostaglandin H2 Synthase 1
Immunogen:	Peptide from the internal region of mouse COX-1
Cross Reactivity:	(-) COX-2
Species Reactivity:	(+) Mouse and ovine; (-) Human; other species not tested
Uniprot No.:	P22437
Form:	Liquid
Storage:	-20°C (as supplied)
Stability:	≥3 years
Storage Buffer:	PBS, pH 7.2, with 50% glycerol and 0.02% sodium azide
Host:	Rabbit
Application:	Western blot; the recommended starting dilution is 1:200. Other applications were not tested, therefore optimal working concentration/dilution should be determined empirically.

Image



Lane 1: Rat brain homogenate (30 µg)

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/15/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM

PRODUCT INFORMATION



Description

Cyclooxygenase 1 (COX-1) is a bifunctional enzyme that exhibits both COX and peroxidase activities.^{1,2} It is composed of an N-terminal signal peptide, an EGF-like domain, a membrane binding domain, a catalytic domain, and a C-terminal tail.³ COX-1 is constitutively expressed in the gastrointestinal tract, kidney, spleen, liver, and lung and localizes to the endoplasmic reticulum.^{4,5} The COX component converts arachidonic acid (Item Nos. 90010 | 90010.1 | 10006607) to a hydroperoxyl endoperoxide prostaglandin G₂ (PGG₂; Item No. 17010) and the peroxidase component reduces the endoperoxide to the corresponding alcohol PGH₂ (Item No. 17020), the precursor of PGs, thromboxanes, and prostacyclins.^{1,2} COX-1 is the target of many non-steroidal anti-inflammatory drugs (NSAIDs) and is responsible for the undesirable gastrointestinal and renal side effects, such as ulcer formation and reductions in the glomerular filtration rate, respectively.^{6,7} Cayman's COX-1 (mouse) Polyclonal Antibody can be used for Western blot (WB).

References

1. Nugteren, D.H. and Hazelhof, E. Isolation and properties of intermediates in prostaglandin biosynthesis. *Biochim. Biophys. Acta* **326(3)**, 448-461 (1973).
2. Hamberg, M. and Samuelsson, B. Detection and isolation of an endoperoxide intermediate in prostaglandin biosynthesis. *Proc. Natl. Acad. Sci. USA* **70(3)**, 899-903 (1973).
3. Smith, W.L., Garavito, R.M., and DeWitt, D.L. Prostaglandin endoperoxide H synthases (cyclooxygenases)-1 and -2. *J. Biol. Chem.* **271**, 33157-33160 (1996).
4. Seibert, K., Zhang, Y., Leahy, K., et al. Pharmacological and biochemical demonstration of the role of cyclooxygenase 2 in inflammation and pain. *Proc. Natl. Acad. Sci. USA* **91(25)**, 12013-12017 (1994).
5. Morita, I., Schindler, M., Regier, M.K., et al. Different intracellular locations for prostaglandin endoperoxide H synthase-1 and -2. *J. Biol. Chem.* **270(18)**, 10902-10908 (1995).
6. Gierse, J.K., Hauser, S.D., Creely, D.P., et al. Expression and selective inhibition of the constitutive and inducible forms of human cyclo-oxygenase. *Biochem. J.* **305(Pt. 2)**, 379-484 (1995).
7. Frölich, J.C. A classification of NSAIDs according to the relative inhibition of cyclooxygenase isoenzymes. *Trends Pharmacol. Sci.* **18(1)**, 30-34 (1997).

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM