



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION




GPR41 (Internal) Polyclonal Antibody

Item No. 16363

Overview and Properties

Contents:	This vial contains 500 µl of peptide affinity-purified polyclonal antibody.
Synonyms:	Free Fatty Acid Receptor 3, G Protein-Coupled Receptor 41
Immunogen:	Synthetic peptide from the internal region of human GPR41
Species Reactivity:	(+) Human; other species not tested
Uniprot No.:	O14843
Form:	Liquid
Storage:	-20°C (as supplied)
Stability:	≥3 years
Storage Buffer:	PBS, pH 7.2, with 50% glycerol and 0.02% sodium azide
Host:	Rabbit
Applications:	ELISA and Western blot (WB); the recommended starting dilution is 1:200. Other applications were not attempted and therefore optimal working dilutions should be determined empirically.

Image



Lane 1: MW Markers
Lane 2: MCF-7 cell lysate (50 µg)
Lane 3: MW Markers
Lane 4: MCF-7 cell lysate (50 µg)
+ 10 µg/ml immunizing
peptide

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/17/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM

PRODUCT INFORMATION



Description

GPR41 is a G protein-coupled receptor activated by short chain fatty acids (SCFAs).¹⁻³ Several SCFAs have the potential to bind and activate GPR41, with pentanoate being the most potent agonist.² GPR41 couples through the Pertussis toxin-sensitive $G_{i/o}$ family and its expression has been described in adipose tissue and the colonic lumen.¹⁻³ The activation of GPR41 induces an increase in intracellular Ca^{2+} , ERK1/2 activation and a decrease in intracellular cAMP.¹⁻³ Activation of GPR41 may be involved in intestinal inflammation. The predicted size for GPR41 is 39 kDa. Cayman's GPR41 (Internal) Polyclonal Antibody detects a 45 kDa band by WB in cell lysates.

References

1. Tazoe, H., Otomo, Y., Kaji, I., *et al.* Roles of short-chain fatty acids receptors, GPR41 and GPR43 on colonic functions. *J. Physiol. Pharmacol.* **59**(Suppl 2), 251-262 (2008).
2. Brown, A.J., Goldsworthy, S.M., Barnes, A.A., *et al.* The orphan G protein-coupled receptors GPR41 and GPR43 are activated by propionate and other short chain carboxylic acids. *J. Biol. Chem.* **278**(13), 11312-11319 (2003).
3. Le Poul, E., Loison, C., Struyf, S., *et al.* Functional characterization of human receptors for short chain fatty acids and their role in polymorphonuclear cell activation. *J. Biol. Chem.* **278**(28), 25481-24591 (2003).

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM