

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



UNC2025

Item No. 16613

CAS Registry No.: 1429881-91-3

Formal Name: trans-4-[2-(butylamino)-5-[4-

> [(4-methyl-1-piperazinyl)methyl] phenyl]-7H-pyrrolo[2,3-d] pyrimidin-7-yl]-cyclohexanol

MF: $C_{28}H_{40}N_6O$ FW: 476.7 **Purity:** ≥98%

UV/Vis.: λ_{max} : 222, 254, 290, 337 nm

A crystalline solid Supplied as:

-20°C Storage:

Stability: As supplied, 2 years from the QC date provided on the Certificate of Analysis, when

stored properly

Laboratory Procedures

UNC2025 is supplied as a crystalline solid. A stock solution may be made by dissolving the UNC2025 in the solvent of choice. UNC2025 is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of UNC2025 in ethanol and DMF is approximately 30 mg/ml, and approximately 10 mg/ml in DMSO.

UNC2025 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, UNC2025 should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. UNC2025 has a solubility of approximately 0.1 mg/ml in a 1:9 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

UNC2025 is a potent, orally bioavailable inhibitor of the tyrosine kinases Mer and Flt3 (IC_{50} s = 0.74 and 0.80 nM, respectively). It less potently inhibits AxI and Tyro3 (IC $_{50}$ s = 14 and 17 nM, respectively) and a panel of related kinases. UNC2025 blocks colony formation of Mer- and Flt3-dependent tumor cell lines and inhibits Mer phosphorylation in bone marrow leukemia cells in vivo. Pharmacokinetic studies suggest that orally administered UNC2025 at 3 mg/kg will suffice to provide 90% inhibition of Mer and Flt3 at 30 min.¹

Reference

1. Zhang, W., DeRychere, D., Hunter, D., et al. UNC2025, a potent and orally bioavailable MER/FLT3 dual inhibitor. J. Med. Chem. 57(16), 7031-7041 (2014).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 06/22/2016

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM