

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



RS 504393

Item No. 17330

CAS Registry No.: 300816-15-3

Formal Name: 6-methyl-1'-[2-(5-methyl-2-phenyl-

4-oxazolyl)ethyl]-spiro[4H-3,1-

benzoxazine-4,4'-piperidin]-2(1H)-one

MF: $C_{25}H_{27}N_3O_3$ 417.5 FW:

≥98% **Purity:**

UV/Vis.: λ_{max} : 246, 281 nm

Supplied as: A solid Storage:

As supplied, 2 years from the QC date provided on the Certificate of Analysis, when Stability:

stored properly

Laboratory Procedures

RS 504393 is supplied as a solid. A stock solution may be made by dissolving the RS 504393 in the solvent of choice. RS 504393 is soluble in organic solvents such as DMSO and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of RS 504393 in DMSO and DMF is approximately 1 and 10 mg/ml, respectively. RS 504393 is also sparingly soluble in ethanol.

RS 504393 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, RS 504393 should first be dissolved in DMF and then diluted with the aqueous buffer of choice. RS 504393 has a solubility of approximately 0.1 mg/ml in a 1:9 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

RS 504393 is a selective antagonist of the MCP-1 receptor CCR2 (IC $_{50}$ s = 89 nM and >100 μ M for CCR2 and CCR1, respectively). It has been shown to inhibit MCP-1-stimulated chemotaxis (IC50 = 330 nM) and ischemia-reperfusion injury in kidneys.^{1,2}

References

- 1. Mirzadegan, T., Diehl, F., Ebi, B., et al. Identification of the binding site for a novel class of CCR2b chemokine receptor antagonists: Binding to a common chemokine receptor motif within the helical bundle. J. Biol. Chem. 275(33), 25562-25571 (2000).
- 2. Furuichi, K., Wada, T., Iwata, Y., et al. CCR2 signaling contributes to ischemia-reperfusion injury in kidney. J. Am. Soc. Nephrol. 14(10), 2503-2515 (2003).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 02/22/2017

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM