



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



PRODUCT INFORMATION



Tasquinimod

Item No. 17692

CAS Registry No.: 254964-60-8

Formal Name: 1,2-dihydro-4-hydroxy-5-methoxy-N,1-dimethyl-2-oxo-N-[4-(trifluoromethyl)phenyl]-3-quinolinecarboxamide

Synonym: ABR-215050

MF: $C_{20}H_{17}F_3N_2O_4$

FW: 406.4


Purity: $\geq 98\%$

UV/Vis.: λ_{max} : 232, 303 nm

Supplied as: A crystalline solid

Storage: $-20^{\circ}C$

Stability: As supplied, 2 years from the QC date provided on the Certificate of Analysis, when stored properly



Laboratory Procedures

Tasquinimod is supplied as a crystalline solid. A stock solution may be made by dissolving the tasquinimod in the solvent of choice. Tasquinimod is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of tasquinimod in these solvents is approximately 2, 10, and 20 mg/ml, respectively.

Tasquinimod is sparingly soluble in aqueous solutions. To enhance aqueous solubility, dilute the organic solvent solution into aqueous buffers or isotonic saline. If performing biological experiments, ensure the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. We do not recommend storing the aqueous solution for more than one day.

Description

Tasquinimod is an orally-active quinoline-3-carboxamide derived from roquinimex, an immunomodulatory quinolone with applications in some cancers and autoimmune diseases.¹ Tasquinimod inhibits tumor angiogenesis and supplements radiation or chemotherapy in animal models of prostate cancer.^{1,2} While its precise mechanism of action remains unclear, tasquinimod has been reported to alter signaling through S100A9, thrombospondin-1, HIF-1 α , androgen receptor, VEGF, and HDAC3/4.^{1,2}

References

1. Isaacs, J.T. The long and winding road for the development of tasquinimod as an oral second-generation quinoline-3-carboxamide antiangiogenic drug for the treatment of prostate cancer. *Expert Opin. Investig. Drugs* **19**(10), 1235-1243 (2010).
2. Mehta, A.R. and Armstrong, A.J. Tasquinimod in the treatment of castrate-resistant prostate cancer - current status and future prospects. *Ther. Adv. Urol.* **8**(1), 9-18 (2016).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 04/19/2016

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM