

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Dynorphin B

Item No. 18178

CAS Registry No.: Formal Name:	83335-41-5 dynorphin B (swine)	
Synonyms:	Dynorphin B-13, Rimorphin	
MF: FW: Purity: Stability: Supplied as: UV/Vis.:	C ₇₄ H ₁₁₅ N ₂₁ O ₁₇ 1,570.8 ≥95% ≥2 years at -20°C A crystalline solid $λ_{max}$: 278 nm	

Laboratory Procedures

For long term storage, we suggest that dynorphin B be stored as supplied at -20°C. It should be stable for at least two years.

Dynorphin B is supplied as a crystalline solid. A stock solution may be made by dissolving the dynorphin B in the solvent of choice. Dynorphin B is soluble in organic solvents such as DMSO and dimethyl formamide, which should be purged with an inert gas. The solubility of dynorphin B in these solvents is approximately 30 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of dynorphin B can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of dynorphin B in PBS, pH 7.2, is approximately 5 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Dynorphin B is a 13-residue opioid peptide released from the cleavage of prodynorphin and found widely distributed in the central nervous system.^{1,2} It acts as a κ_{1b} -opioid receptor agonist (K_i = 1.1 nM) and has been implicated in antinociceptive functions.³ Dynorphin B has also been used to prime cardiogenesis in pluripotent embryonic stem cells.4

References

- 1. Berman, Y., Juliano, L., and Devi, L.A. Specificity of the dynorphin-processing endoprotease: Comparison with prohormone convertases. J. Neurochem. 72, 2120-2196 (1999).
- 2. Janecka, A., Fichna, J., and Janecki, T. Opioid receptors and their ligands. Curr. Top. Med. Chem. 4, 1-17 (2004).
- 3. Dhawan, B.N., Cesselin, F., Raghubir, R., et al. International Union of Pharmacology. XII. Classification of opioid receptors. Pharm. Res. 48(4), 567-592 (1996).
- 4. Ventura, C., Zinellu, E., Maninchedda, E., et al. Dynorphin B is an agonist of nuclear opioid receptors coupling nuclear protein kinase C activation to the transcription of cardiogenic genes in GTR1 embryonic stem cells. Circ. Res. 92, 623-629 (2003).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

al should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/10/2015

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM