

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



RS 102895 (hydrochloride)

Item No. 18207

| CAS Registry No.: Formal Name: | 1173022-16-6 1'-[2-[4-(trifluoromethyl)phenyl]ethyl]- spiro[4H-3,1-benzoxazine-4,4'-piperidin]- 2(1H)-one, monohydrochloride | |
|-----------------------------------|---|-------|
| MF: | $C_{21}H_{21}F_3N_2O_2 \bullet HCI$ | • HCI |
| FW: | 426.9 | |
| Purity: | ≥98% | |
| Stability: | ≥2 years at -20°C | |
| Supplied as: | A crystalline solid | ~ N - |
| UV/Vis.: | λ _{max} : 204, 279 nm | Ĥ |

Laboratory Procedures

For long term storage, we suggest that RS 102895 (hydrochloride) be stored as supplied at -20°C. It should be stable for at least two years.

RS 102895 (hydrochloride) is supplied as a crystalline solid. A stock solution may be made by dissolving the RS 102895 (hydrochloride) in the solvent of choice. RS 102895 (hydrochloride) is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of RS 102895 (hydrochloride) in these solvents is approximately 2, 30, and 25 mg/ml, respectively.

RS 102895 (hydrochloride) is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, RS 102895 (hydrochloride) should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. RS 102895 (hydrochloride) has a solubility of approximately 0.3 mg/ml in a 1:2 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

The chemokine CCL2, also known as monocyte chemotactic protein-1 (MCP-1), stimulates leukocyte chemotaxis to sites of inflammation via signaling through the MCP-1 receptor, CCR2. RS 102895 is a spiropiperidine compound that acts as a CCR2 antagonist (IC₅₀ = 0.36 μ M).¹ It inhibits the related CCR1 receptor with an IC₅₀ value of 17.8 μ M and inhibits adrenergic α_{1a} , α_{1d} , and 5HT_{1A} receptors with IC₅₀ values of 0.13, 0.32, and 47 μ M, respectively.¹ RS 102895 prevents chemotaxis of THP-1 cells $(IC_{50} = 1.7 \,\mu\text{M})$ when MCP-1 is presented as a chemoattractant.¹

Reference

1. Mirzadegan, T., Diehl, F., Ebi, B., et al. Identification of the binding site for a novel class of CCR2b chemokine receptor antagonists: Binding to a common chemokine receptor motif within the helical bundle. J. Biol. Chem. 275(33), 25562-25571 (2000).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 09/08/2015

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM