



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PRODUCT INFORMATION



COX-2 (ovine, recombinant)


Item No. 18356

Overview and Properties

Synonyms:	Cyclooxygenase 2, Inducible Cyclooxygenase Prostaglandin H Synthase 2
Source:	Active recombinant ovine COX-2 expressed in insect cells
Amino Acids:	24-599
Uniprot No.:	P79208
Molecular Weight:	~70 kDa
Storage:	-80°C (as supplied); avoid freeze/thaw cycles by aliquoting protein
Stability:	≥6 months
Purity:	batch specific (≥80% estimated by SDS-PAGE)
Supplied in:	batch specific
Activity:	batch specific U/ml. One unit is defined as the amount of enzyme required to consume 1 nmol of oxygen per minute 37°C in 100 mM Tris, pH 8.0, containing 100 µM arachidonate, 5 mM EDTA, 2 mM phenol, and 1 µM hematin.
Specific Activity:	batch specific nmol/min/mg
Protein Concentration:	batch specific mg/ml
Specific Activity:	>20,000 U/mg

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Image



Lane 1: MW Markers
Lane 2: COX-2 (2 µg)
Lane 3: COX-2 (5 µg)

SDS-PAGE Analysis of COX-2.

Representative gel image shown; actual purity may vary between each batch.

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 06/04/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 - USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM

PRODUCT INFORMATION

Description

Cyclooxygenase 2 (COX-2) is a bifunctional enzyme that exhibits both COX and peroxidase activities and catalyzes the first step in the biosynthesis of prostaglandins, thromboxanes, and prostacyclins.^{1,2} The COX component converts arachidonic acid to the hydroperoxy endoperoxide prostaglandin G₂ (PGG₂; Item No. 17010), and the peroxidase component reduces the endoperoxide to the corresponding alcohol PGH₂ (Item No. 17020). COX2 expression is induced by a variety of stimuli, including phorbol esters, LPS, and cytokines and is responsible for the biosynthesis of PGs under acute inflammatory conditions.^{3,4} Thus, COX-2 has been the focus of attention for nonsteroidal anti-inflammatory drug (NSAID) development. Cayman's COX-2 (ovine, recombinant) protein can be used for enzyme activity assays.

References

1. Nugteren, D.H. and Hazelhof, E. Isolation and properties of intermediates in prostaglandin biosynthesis. *Biochim. Biophys. Acta* **326**(3), 448-461 (1973).
2. Hamberg, M. and Samuelsson, B. Detection and isolation of an endoperoxide intermediate in prostaglandin biosynthesis. *Proc. Natl. Acad. Sci. USA* **70**(3), 899-903 (1973).
3. Kang, Y.-J., Mbonye, U.R., DeLong, C.J., et al. Regulation of intracellular cyclooxygenase levels by gene transcription and protein degradation. *Prog. Lipid Res.* **46**(2), 108-25 (2007).
4. Blobaum, A.L. and Marnett, L.J. Structural and functional basis of cyclooxygenase inhibition. *J. Med. Chem.* **50**(7), 1425-1441 (2007).