

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Iopromide

Item No. 18470

CAS Registry No.: 73334-07-3

Formal Name: N¹,N³-bis(2,3-dihydroxypropyl)-

2,4,6-triiodo-5-[(2-methoxyacetyl)

amino]-N¹-methyl-1,3-

benzenedicarboxamide

Synonym: Ultravist $C_{18}H_{24}I_3N_3O_8$ MF:

791.1 FW: **Purity:** ≥98%

Stability: ≥2 years at -20°C Supplied as: A crystalline solid UV/Vis.: λ_{max} : 241 nm

Laboratory Procedures

For long term storage, we suggest that iopromide be stored as supplied at -20°C. It should be stable for at least two years.

lopromide is supplied as a crystalline solid. A stock solution may be made by dissolving the iopromide in the solvent of choice. lopromide is soluble in organic solvents such as DMSO and dimethyl formamide, which should be purged with an inert gas. The solubility of iopromide in these solvents is approximately 10 and 3 mg/ml, respectively.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of iopromide can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of iopromide in PBS, pH 7.2, is approximately 3 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

lopromide is a non-ionic, water-soluble X-ray contrast agent for intravascular administration.^{1,2} lopromide can cause acute kidney injury and is used in animal models of contrast-induced nephropathy.^{3,4}

References

- 1. Ahn, Y.-H., Koh, Y.-I., Kim, J.-H., et al. The potential utility of iodinated contrast media (ICM) skin testing in patients with ICM hypersensitivity. J. Korean Med. Sci. 30(3), 245-251 (2015).
- Park, J.-Y., Lee, S.-K., Kim, J.-Y., et al. A new micro-computed tomography-based high-resolution bloodbrain barrier imaging technique to study ischemic stroke. Stroke 45(8), 2480-2484 (2014).
- Hassan, K. and Fadi, H. Is hypoalbuminemia a prognostic risk factor for contrast-induced nephropathy in peritoneal dialysis patients? Ther. Clin. Risk Manag. 10, 787-795 (2014).
- Wang, N., Wei, R., Li, Q., et al. Renal Protective Effect of Probucol in Rats with Contrast-Induced Nephropathy and its Underlying Mechanism. Med. Sci. Monit. 21, 2886-2892 (2015).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/17/2015

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA **PHONE:** [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM