

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



ABT-333

Item No. 18482

CAS Registry No.: 1132935-63-7

Formal Name: N-[6-[5-(3,4-dihydro-2,4-

> dioxo-1(2H)-pyrimidinyl)-3-(1,1-dimethylethyl)-2-

methoxyphenyl]-2-naphthalenyl]-

methanesulfonamide

Synonym: Dasabuvir MF: $C_{26}H_{27}N_3O_5S$ FW: 493.5 **Purity:**

UV/Vis.: λ_{max} : 244 nm Supplied as: A crystalline solid

-20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



ABT-333 is supplied as a crystalline solid. A stock solution may be made by dissolving the ABT-333 in the solvent of choice. ABT-333 is soluble in organic solvents such as DMSO and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of ABT-333 in these solvents is approximately 20 and 30 mg/ml, respectively.

ABT-333 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, ABT-333 should first be dissolved in DMF and then diluted with the aqueous buffer of choice. ABT-333 has a solubility of approximately 0.25 mg/ml in a 1:3 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

ABT-333 is an inhibitor of hepatitis C virus (HCV) RNA-dependent RNA polymerase (RdRp; IC₅₀s = 0.0022-0.0107 μM for genotypes 1a and 1b).¹ It is selective for HCV genotype 1 polymerases over genotypes 2, 3, and 4 (IC_{50} s = 0.9->20 μ M), as well as human polymerases (IC_{50} s = >100 μ M). Formulations containing ABT-333, ombitasvir, paritaprevir, and ritonavir have been used in the treatment of HCV genotype 1 infections.

Reference

1. Kati, W., Koev, G., Irvin, M., et al. In vitro activity and resistance profile of dasabuvir, a nonnucleoside hepatitis C virus polymerase inhibitor. Antimicrob. Agents Chemother. 59(3), 1505-1511 (2015).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/08/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM