

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



BPTFS

Item No. 19284

CAS Registry No.: 314045-39-1

Formal Name: N,N'-[thiobis(2,1-ethanediyl-

1,3,4-thiadiazole-5,2-diyl)]bis-


benzeneacetamide

MF: $C_{24}H_{24}N_6O_2S_3$

FW: 524.7 ≥95% **Purity:** UV/Vis.: λ_{max} : 254 nm A crystalline solid Supplied as:

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

BPTES is supplied as a crystalline solid. A stock solution may be made by dissolving the BPTES in the solvent of choice. BPTES is soluble in organic solvents such as DMSO and dimethyl formamide, which should be purged with an inert gas. The solubility of BPTES in these solvents is approximately 20 and 10 mg/ml, respectively.

BPTES is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, BPTES should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. BPTES has a solubility of approximately 0.33 mg/ml in a 1:2 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

BPTES is an allosteric inhibitor of kidney-type glutaminase 1 (GLS1; IC $_{50}$ = 3.3 μ M) that drives the formation of inactive GLS1 tetramers. It is selective for GLS1 over GLS2, glutamate dehydrogenase, and y-glutamyl transpeptidase. BPTES decreases glutaminase activity and reduces proliferation of aerobic P493 cells as well as induces cell death, increases the production of reactive oxygen species (ROS), and reduces ATP levels in hypoxic P493 cells.² It also decreases ATP-linked and uncoupled oxygen consumption in C2C12 myotubes when used in combination with UK 5099 (Item No. 16980) and/or etomoxir (Item No. 11969).³ In vivo, BPTES (12.5 mg/kg) reduces tumor volume in a P493 lymphoma mouse xenograft model.²

References

- 1. Shukla, K., Feraris, D.V., Thomas, A.G., et al. Design, synthesis, and pharmacological evaluation of bis-2-(5-phenylacetamido-1,2,4-thiadiazol-2-yl)ethyl sulfide 3 (BPTES) analogs as glutaminase inhibitors. J. Med. Chem. 55(23), 10551-10563 (2012).
- 2. Le, A., Lane, A.N., Hamaker, M., et al. Glucose-independent glutamine metabolism via TCA cycling for proliferation and survival in B cells. Cell. Metab. 15(1), 110-121 (2012).
- Vacanti, N.M., Divakaruni, A.S., Green, C.R., et al. Regulation of substrate utilization by the mitochondrial pyruvate carrier. Mol. Cell 56(3), 425-435 (2014).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 12/19/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM