



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



Eletriptan

Item No. 20048

CAS Registry No.: 143322-58-1
Formal Name: 3-[[[(2R)-1-methyl-2-pyrrolidinyl]methyl]-5-[2-(phenylsulfonyl)ethyl]-1H-indole
Synonym: UK 116044
MF: C₂₂H₂₆N₂O₂S
FW: 382.5
Purity: ≥98%
UV/Vis.: λ_{max}: 223, 271 nm
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Eletriptan is supplied as a crystalline solid. A stock solution may be made by dissolving the eletriptan in the solvent of choice. Eletriptan is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of eletriptan in ethanol is approximately 0.1 mg/ml, and approximately 10 mg/ml in DMSO and DMF.

Eletriptan is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, eletriptan should first be dissolved in DMF and then diluted with the aqueous buffer of choice. Eletriptan has a solubility of approximately 0.5 mg/ml in a 1:1 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Eletriptan is an agonist of the serotonin (5-HT) receptor types 5-HT_{1B} and 5-HT_{1D}.¹ It inhibits forskolin-induced cAMP accumulation in and induces vasoconstriction of isolated rabbit common carotid artery rings (pD₂s = 7.6 and 4.9, respectively), an effect that can be blocked by the 5-HT_{1B} antagonist SB216641 but not the 5-HT_{1D} antagonist BRL15572. Eletriptan preferentially induces constriction of isolated human cerebral over coronary arteries (EC₅₀s = 15.8 and 1,995 nM, respectively).² Formulations containing eletriptan have been used in the treatment of migraine headache.

References

1. Akin, D., Onaran, H. O., and Gudal, H. Agonist-directed trafficking explaining the difference between response pattern of naratriptan and sumatriptan in rabbit common carotid artery. *Br. J. Pharmacol.* **136**(2), 171-176 (2002).
2. Edvinsson, L., Uddman, E., Wackenfors, A., *et al.* Triptan-induced contractile (5-HT_{1B} receptor) responses in human cerebral and coronary arteries: Relationship to clinical effect. *Clin. Sci. (Lond.)* **109**(3), 335-342 (2005).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 06/20/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM