

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



EMPO

Item No. 20618

CAS Registry No.: 61856-99-3

Formal Name: 3,4-dihydro-2-methyl-2H-Pyrrole-2-carboxylic acid

1-oxide, ethyl ester

Synonym: 5-(Ethoxycarbonyl)-5-methyl-1-Pyrroline-N-Oxide

MF: $C_8H_{13}NO_3$ 171.2 FW: ≥98% **Purity:** UV/Vis.: λ_{max} : 238 nm Supplied as: A solution in ethanol

Storage:

As supplied, 2 years from the QC date provided on the Certificate of Analysis, when Stability:

stored properly



EMPO is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide (DMF) purged with an inert gas can be used. The solubility of EMPO in ethanol and DMF is approximately 30 mg/ml and approximately 15 mg/ml in DMSO.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of EMPO is needed, it can be prepared by evaporating the ethanol and directly dissolving the neat oil in aqueous buffers. The solubility of EMPO in PBS, pH 7.2, is approximately 1 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

EMPO is a hydrophilic cyclic nitrone analog of the free radical spin trap DMPO (Item No. 10006436). The EMPO-superoxide adduct (EMPO-OOH) exhibits an ESR spectrum that is similar to the DMPO-OOH spectrum. However, the EMPO-OOH adduct is about 5-8 times more stable ($t_{1/2} = 4.8 - 8.6$ min) and does not spontaneously decay to the hydroxyl adduct, making spectral interpretation less difficult.¹⁻³

References

- 1. Olive, G., Mercier, A., Le Moigne, F., et al. 2-ethoxycarbonyl-2-methyl-3,4-dihydro-2H-pyrrole-1-oxide: Evaluation of the spin trapping properties. Free Radic. Biol. Med. 28(3), 403-408 (2000).
- 2. Stolze, K., Udilova, N., and Nohl, H. Spin adducts of superoxide, alkoxyl, and lipid-derived radicals with EMPO and its derivatives. Biol. Chem. 383(5), 813-820 (2002).
- Zhang, H., Joseph, J., Vasquez-Vivar, J., et al. Detection of superoxide anion using an isotopically labeled nitrone spin trap: Potential biological applications. FEBS Lett. 473(1), 58-62 (2000).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 01/04/2017

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM