

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Chlorprothixene (hydrochloride)

Item No. 20772

CAS Registry No.: 6469-93-8

Formal Name: (3Z)-3-(2-chloro-9H-thioxanthen-9-ylidene)-N,N-

dimethyl-1-propanamine, monohydrochloride

Synonyms: cis-Chlorprothixene, NSC 169899

C₁₈H₁₈CINS • HCI MF:

352.3 FW: **Purity:** ≥98%

UV/Vis.: λ_{max} : 230, 270, 329 nm Supplied as: A crystalline solid

Storage:

As supplied, 2 years from the QC date provided on the Certificate of Analysis, when Stability:

stored properly



Chlorprothixene (hydrochloride) is supplied as a crystalline solid. A stock solution may be made by dissolving the chlorprothixene (hydrochloride) in the solvent of choice. Chlorprothixene (hydrochloride) is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of chlorprothixene (hydrochloride) in these solvents is approximately 30 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of chlorprothixene (hydrochloride) can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of chlorprothixene (hydrochloride) in PBS, pH 7.2, is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Chlorprothixene is a thioxanthine antipsychotic that functions by antagonizing dopamine D_2 receptors. It can block a subset of GABA_∆ receptors in rat cortex that is also blocked by clozapine (Item No. 12059).² It has also been shown to be effective against P. falciparum growth with an EC₅₀ value of 1.7 μ M.³

References

- 1. Froimowitz, M., and Cody, V. Biologically active conformers of phenothiazines and thioxanthenes. Further evidence for a ligand model of dopamine D₂ receptor antagonists. J. Med. Chem. 36(15), 2219-2227
- 2. Squires, R.F., and Saederup, E. Clozapine and several other antipsychotic/antidepressant drugs preferentially block the same 'core' fraction of GABA, receptors. Neurochem. Res. 23(10), 1283-1290
- 3. Weisman, J.L., Liou, A.P., Shelat, A.A., et al. Searching for new antimalarial therapeutics amongst known drugs. Chem. Biol. Drug Des. 67(6), 409-416 (2006).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 12/07/2016

• HCI

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM