



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



2,3-Butanedione-2-monoxime

Item No. 20828

CAS Registry No.:	57-71-6
Formal Name:	2,3-butanedione, 2-oxime
Synonyms:	BDM, Diacetyl monoxime, NSC 660, NSC 116103
MF:	C ₄ H ₇ NO ₂
FW:	101.1
Purity:	≥95%
UV/Vis.:	λ _{max} : 229 nm
Supplied as:	A crystalline solid
Storage:	-20°C
Stability:	As supplied, 2 years from the QC date provided on the Certificate of Analysis, when stored properly



Laboratory Procedures

2,3-Butanedione-2-monoxime (BDM) is supplied as a crystalline solid. A stock solution may be made by dissolving the BDM in the solvent of choice. BDM is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of BDM in these solvents is approximately 30 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of BDM can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of BDM in PBS, pH 7.2, is approximately 2 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

BDM is a non-selective, reversible myosin ATPase inhibitor with phosphatase-like properties.^{1,2} It demonstrates direct and indirect actions on cardiac and vascular smooth muscle, inhibiting L-type Ca²⁺ channels.^{3,4} BDM has been shown to be involved in a number of biological activities, including muscle contraction and synaptic transmission.¹

References

1. Sellin, L.C. and McArdle, J.J. Multiple effects of 2,3-butanedione monoxime. *Pharmacol. Toxicol.* **74**(6), 305-313 (1994).
2. Ostap, E.M. 2,3-Butanedione monoxime (BDM) as a myosin inhibitor. *J. Muscle Res. Cell Motil.* **23**(4), 305-308 (2002).
3. Eisfeld, J., Mikala, G., Varadi, G., *et al.* Inhibition of cloned human L-type cardiac calcium channels by 2,3-butanedione monoxime does not require PKA-dependent phosphorylation sites. *Biochem. Biophys. Res. Commun.* **230**(3), 489-492 (1997).
4. Adhikari, B.B. and Wang, K. Interplay of troponin- and myosin-based pathways of calcium activation in skeletal and cardiac muscle: The use of W7 as an inhibitor of thin filament activation. *Biophys. J.* **86**(1), 359-370 (2004).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/05/2016

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM