

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Etodolac

Item No. 20833

CAS Registry No.: 41340-25-4

Formal Name: 1,8-diethyl-1,3,4,9-tetrahydro-

pyrano[3,4-b]indole-1-acetic acid

Synonyms: AY 24236, (±)-Etodolac, NIH 9918,

NSC 282126

MF: $C_{17}H_{21}NO_3$ FW: 287.4 **Purity:** ≥98%

UV/Vis.: λ_{max} : 224, 273 nm Supplied as: A crystalline solid Storage: Room temperature

Stability: As supplied, 2 years from the QC date provided on the Certificate of Analysis, when

stored properly

Laboratory Procedures

Etodolac is supplied as a crystalline solid. A stock solution may be made by dissolving the etodolac in the solvent of choice. Etodolac is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of etodolac in ethanol is approximately 20 mg/ml and approximately 30 mg/ml in DMSO and DMF.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of etodolac can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of etodolac in PBS, pH 7.2, is approximately 0.25 mg/ml. We do not recommend storing the agueous solution for more than one day.

Description

Etodolac is a non-steroidal anti-inflammatory drug (NSAID) that inhibits both COX isoforms in vitro, with modest selectivity for COX-2 (IC₅₀s = 12 μ M for COX-1 and 2.2 μ M for COX-2 in a human whole blood assay).1 Experimental evidence suggests that etodolac may have favorable tissue pharmacokinetics resulting in further COX-2 selectivity, indicated by favorable reduction in anti-inflammatory action with diminished gastric side effects.^{2,3}

References

- 1. Warner, T.D., Giuliano, F., Vojnovic, I., et al. Nonsteroid drug selectivities for cyclo-oxygenase-1 rather than cyclo-oxygenase-2 are associated with human gastrointestinal toxicity: A full in vitro analysis. Proc. Natl. Acad. Sci. USA 96, 7563-7568 (1999).
- 2. Dvornik, D.M. Tissue selective inhibition of prostaglandin biosynthesis by etodolac. J. Rheumatol. 24, 40-50 (1997).
- 3. Riendeau, D., Percival, M.D., Brideau, C., et al. Etoricoxib (MK-0663): Preclinical profile and comparison with other agents that selectively inhibit cyclooxygenase-2. J. Pharmacol. Exp. Ther. 296, 558-566 (2001).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/14/2016

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA **PHONE:** [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM