



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



DMAT

Item No. 21182

CAS Registry No.: 749234-11-5

Formal Name: 4,5,6,7-tetrabromo-N,N-dimethyl-1H-benzimidazol-2-amine

Synonyms: Casein Kinase II Inhibitor II,
2-Dimethylamino-4,5,6,7-tetrabromobenzimidazole

MF: $C_9H_7Br_4N_3$

FW: 476.8


Purity: $\geq 98\%$

UV/Vis.: λ_{max} : 239, 271, 315 nm

Supplied as: A crystalline solid

Storage: -20°C

Stability: As supplied, 2 years from the QC date provided on the Certificate of Analysis, when stored properly



Laboratory Procedures

DMAT is supplied as a crystalline solid. A stock solution may be made by dissolving the DMAT in the solvent of choice. DMAT is soluble in organic solvents such as DMSO and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of DMAT in these solvents is approximately 20 and 30 mg/ml, respectively.

DMAT is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, DMAT should first be dissolved in DMF and then diluted with the aqueous buffer of choice. DMAT has a solubility of approximately 0.20 mg/ml in a 1:4 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

DMAT is a cell-permeable inhibitor of casein kinase 2 (CK2; $IC_{50} = 0.13 \mu\text{M}$).¹ It also inhibits Pim-1, Pim-3, HIPK2, and HIPK3 (IC_{50} s = 0.15, 0.097, 0.37, and 0.59 μM , respectively).¹ DMAT blocks cell growth and induces cell death in cancer cells, both in culture and in mouse xenografts.²⁻⁴

References

1. Pagano, M.A., Bain, J., Kazimierczuk, Z., *et al.* The selectivity of inhibitors of protein kinase CK2: An update. *Biochem. J.* **415**(3), 353-365 (2008).
2. Lawnicka, H., Kowalewicz-Kulbat, M., Sicinska, P., *et al.* Anti-neoplastic effect of protein kinase CK2 inhibitor, 2-dimethylamino-4,5,6,7-tetrabromobenzimidazole (DMAT), on growth and hormonal activity of human adrenocortical carcinoma cell line (H295R) *in vitro*. *Cell Tissue Res.* **340**(2), 371-379 (2010).
3. Sass, G., Klinger, N., Surma, H., *et al.* Inhibition of experimental HCC growth in mice by use of the kinase inhibitor DMAT. *Int. J. Oncol.* **39**(2), 433-442 (2011).
4. Yde, C.W., Frogne, T., Lykkesfeldt, A.E., *et al.* Induction of cell death in antiestrogen resistant human breast cancer cells by the protein kinase CK2 inhibitor DMAT. *Cancer Lett.* **256**(2), 229-237 (2007).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 04/10/2017

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM