

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Pyropheophorbide *a*

Item No. 21371

CAS Registry No.: Formal Name:	24533-72-0 9-ethenyl-14-ethyl-4S,8,13,18- tetramethyl-20-oxo-3S- phorbinepropanoic acid	O OH
MF:	$C_{33}H_{34}N_4O_3$	
FW:	534.7	
Purity:	≥95%	
UV/Vis.:	λ _{max} : 209, 226, 273, 324, 409, 507, 539, 609, 666 nm	
Supplied as:	A crystalline solid	
Storage:	-20°C	
Stability:	≥4 years	
Item Origin:	Natural/Source unknown	

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Pyropheophorbide a is supplied as a crystalline solid. A stock solution may be made by dissolving the pyropheophorbide a in the solvent of choice, which should be purged with an inert gas. Pyropheophorbide a is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of pyropheophorbide a in these solvents is approximately 1 mg/ml.

Pyropheophorbide a is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, pyropheophorbide a should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. Pyropheophorbide a has a solubility of approximately 0.25 mg/ml in a 1:3 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Pyropheophorbide a is a photosensitizer and degradant of chlorophyll a (Item No. 33399) that has been found in O. ficus indica and has anticancer and antiviral activities.¹⁻³ It induces cytotoxicity in SKOV3 ovarian cancer and KB papilloma cells when combined with photodynamic therapy (PDT; $IC_{so}s = 14.5$ and 1.2 μ M, respectively).¹ Pyropheophorbide *a* also inhibits the replication of herpes simplex virus 2 (HSV-2) and influenza A virus in Vero and MDCK cells, respectively (IC₅₀s = 0.016 and 0.33 μ g/ml, respectively).²

Reference

- 1. Stamati, I., Kuimova, M.K., Lion, M., et al. Novel photosensitisers derived from pyropheophorbide-a: Uptake by cells and photodynamic efficiency in vitro. Photochem. Photobiol. Sci. 9(7), 1033-1041 (2010).
- 2. Bouslama, L., Hayashi, K., Lee, J.-B., et al. Potent virucidal effect of pheophorbide a and pyropheophorbide a on enveloped viruses. J. Nat. Med. 65(1), 229-233 (2011).
- 3. Doi, M., Inage, T., and Shioi, Y. Chlorophyll degradation in a Chlamydomonas reinhardtii mutant: An accumulation of pyropheophorbide a by anaerobiosis. Plant Cell Physiol. 42(5), 469-474 (2001).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/25/2025

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM