

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Tolnaftate

Item No. 21446

CAS Registry No.: 2398-96-1

Formal Name: N-methyl-N-(3-methylphenyl)-

carbamothioic acid, O-2-naphthalenyl ester

Synonyms: NSC 233648, SCH 10144

MF: C₁₉H₁₇NOS 307.4 FW: **Purity:** ≥98%

UV/Vis.: λ_{max} : 223, 258 nm Supplied as: A crystalline solid

Storage:

As supplied, 2 years from the QC date provided on the Certificate of Analysis, when Stability:

stored properly

Laboratory Procedures

Tolnaftate is supplied as a crystalline solid. A stock solution may be made by dissolving the tolnaftate in the solvent of choice. Tolnaftate is soluble in organic solvents such as DMSO and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of tolnaftate in these solvents is approximately 15 and 30 mg/ml, respectively. Tolnaftate is also slightly soluble in ethanol.

Tolnaftate is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, tolnaftate should first be dissolved in DMF and then diluted with the aqueous buffer of choice. Tolnaftate has a solubility of approximately 0.33 mg/ml in a 1:2 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Tolnaftate is a thiocarbamate antifungal agent.¹ It is active against clinical isolates of the dermatophytes T. rubrum, T. mentagraphytes, T. verrucosum, E. floccosum, and M. canis (MIC₅₀s = 50, 100, 6, 50, and 50 ng/ml, respectively), as well as 18 additional yeast and filamentous fungi species (MIC = 0.003-0.8 μg/ml).^{2,3} Tolnaftate also reduces aflatoxin production in A. parasiticus in a concentration-dependent manner.⁴ It inhibits squalene epoxidase with an IC_{50} value of 12.5 μ g/ml in a cell-free assay.¹

References

- 1. Ryder, N.S., Frank, I., and Dupont, M.C. Ergosterol biosynthesis inhibition by the thiocarbamate antifungal agents tolnaftate and tolciclate. Antimicrob. Agents Chemother. 29(5), 858-860 (1986).
- Petranyi, G., Meingassner, J.G., and Mieth, H. Antifungal activity of the allylamine derivative terbinafine in vitro. Antimicrob. Agents Chemother. **31(9)**, 1365-1368 (1987).
- Waitz, J.A., Moss, E.L., and Weinstein, M.J. Chemotherapeutic evaluation of clotrimazole [Bay b 5097, 1 (o-chloro-α-α-diphenylbenzyl) imidazole]. Appl. Microbiol. 22(5), 891-898 (1971).
- 4. Khan, S.N., Maggon, K.K., and Venkitasubramanian, T.A. Inhibition of aflatoxin biosynthesis by tolnaftate. Appl. Environ. Microbiol. 36(2), 270-273 (1978).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 01/09/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM