

# Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

# **PRODUCT** INFORMATION



#### M71

Item No. 21622

#### Sold under license from The University of Dundee

CAS Registry No.: Formal Name:	1797406-69-9 (4R)-N-[14-[(6S)-4-(4-chlorophenyl)-2,3,9-trimethyl- 6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6- yl]-1,13-dioxo-3,6,9-trioxa-12-azatetradec-1-yl]-3- methyl-L-valyl-4-hydroxy-N-[[4-(4-methyl-5-thiazolyl) phenyl]methyl]-L-prolinamide		
MF:	$C_{A0}H_{A0}CIN_{9}O_{8}S_{2}$		
FW:	1,002.6	` <_>	
Purity:	≥95%		
UV/Vis.:	λ <sub>may</sub> : 256 nm	G	)s
Supplied as:	A crystalline solid		
Storage:	-20°C		
Stability:	≥4 years		
1 6 1			

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

#### Laboratory Procedures

MZ1 is supplied as a crystalline solid. A stock solution may be made by dissolving the MZ1 in the solvent of choice, which should be purged with an inert gas. MZ1 is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of MZ1 in these solvents is approximately 30 mg/ml.

MZ1 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, MZ1 should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. MZ1 has a solubility of approximately 0.12 mg/ml in a 1:7 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

#### Description

MZ1 is a hybrid compound that drives the selective proteasomal degradation of bromodomain-containing protein 4 (BRD4).<sup>1</sup> It is characterized as a proteolysis-targeting chimera (PROTAC) and contains JQ-1, which binds bromo- and extra-terminal (BET) proteins, linked to a ligand for the E3 ubiguitin ligase VHL. MZ1 induces the selective degradation of BRD4 in HeLa cells when used at 0.1-0.5  $\mu$ M.<sup>1</sup> At concentrations of 2-10 μM, MZ1 causes the removal of BRD2, BRD3, and BRD4.<sup>1</sup> BRD protein removal is evident within four hours of adding MZ1 and persists for at least 48 hours, unless MZ1 is removed.<sup>1</sup>

#### Reference

1. Zengerle, M., Chan, K.-H., and Ciulli, A. Selective small molecule induced degradation of the BET bromodomain protein BRD4. ACS Chem. Biol. 10(8), 1770-1777 (2015).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/23/2022

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM