



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



GLPG0187

Item No. 21792

CAS Registry No.: 1320346-97-1
Formal Name: 3-[[2,5-dimethyl-6-[4-(5,6,7,8-tetrahydro-1,8-naphthyridin-2-yl)-1-piperidinyl]-4-pyrimidinyl]amino]-N-[(4-methoxyphenyl)sulfonyl]-L-alanine
MF: C₂₉H₃₇N₇O₅S
FW: 595.7
Purity: ≥98%
UV/Vis.: λ_{max}: 240 nm
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

GLPG0187 is supplied as a crystalline solid. A stock solution may be made by dissolving the GLPG0187 in the solvent of choice, which should be purged with an inert gas. GLPG0187 is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of GLPG0187 in these solvents is approximately 3 and 1 mg/ml, respectively.

GLPG0187 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, GLPG0187 should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. GLPG0187 has a solubility of approximately 0.5 mg/ml in a 1:1 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

GLPG0187 is a nonpeptide antagonist of α_v integrin receptors (IC₅₀s = 1.2-3.7 nM).¹ It inhibits osteoclastogenesis, bone resorption, and angiogenesis *in vitro* and *in vivo* in mice, which are all hallmarks of bone marrow invasion in metastatic prostate cancer. GLPG0187 reduces epithelial-mesenchymal transition (EMT) and migration of PC-3M-Pro4/luc cells in a dose-dependent manner without inhibiting cell growth. GLPG0187 also induces detachment and necrosis of GL-261 and SMA-560 murine glioma cells.²

References

1. van der Horst, G., van den Hoogen, C., Buijjs, J.T., *et al.* Targeting of α_v-integrins in stem/progenitor cells and supportive microenvironment impairs bone metastasis in human prostate cancer. *Neoplasia* **13**(6), 516-525 (2011).
2. Silginer, M., Weller, M., Ziegler, U., *et al.* Integrin inhibition promotes atypical anoikis in glioma cells. *Cell Death Dis.* **5**, e1012 (2014).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/14/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM