

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# PRODUCT INFORMATION



## preQ<sub>1</sub> (hydrochloride)

Item No. 21864

CAS Registry No.: 86694-45-3

Formal Name: 2-amino-5-(aminomethyl)-3,7-dihydro-4H-

pyrrolo[2,3-d]pyrimidin-4-one, dihydrochloride

Synonym: Pre-Queuosine 1


MF: C7H9N5O • 2HCI

FW: 252.1 **Purity:** ≥90%

UV/Vis.:  $\lambda_{\text{max}}$ : 217, 254 nm Supplied as: A crystalline solid

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



#### **Laboratory Procedures**

preQ<sub>1</sub> (hydrochloride) is supplied as a crystalline solid. A stock solution may be made by dissolving the preQ<sub>1</sub> (hydrochloride) in the solvent of choice, which should be purged with an inert gas. preQ<sub>1</sub> (hydrochloride) is soluble in the organic solvent DMSO at a concentration of approximately 10 mg/ml. Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of preQ1 (hydrochloride) can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of preQ<sub>1</sub> (hydrochloride) in PBS, pH 7.2, is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

#### Description

preQ<sub>1</sub> is a precursor in the biosynthesis of queuosine, a bacterial nucleoside incorporated into tRNA by tRNA transglycosylase. 1 preQ<sub>1</sub> binds to an aptamer domain on tRNA that acts as a riboswitch to repress gene expression.2

#### References

- 1. Okada, N., Noguchi, S., Kasai, H., et al. Novel mechanism of post-transcriptional modification of tRNA. Insertion of bases of Q precursors into tRNA by a specific tRNA transglycosylase reaction. J. Biol. Chem. 254(8), 3067-3073 (1979).
- 2. Wu, M.-C., Lowe, P.T., Robinson, C.J., et al. Rational re-engineering of a transcriptional silencing PreQ1 riboswitch. J. Am. Chem. Soc. 137(28), 9012-9021 (2015).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

### WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 09/07/2021

#### **CAYMAN CHEMICAL**

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM