

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Bicalutamide-d₄

Item No. 22377

CAS Registry No.: 1185035-71-5

Formal Name: N-[4-cyano-3-(trifluoromethyl)phenyl]-

3-[(4-fluorophenyl-2,3,5,6-d₄)sulfonyl]-

2-hydroxy-2-methyl-propanamide

MF: $C_{18}H_{10}D_4F_4N_2O_4S$

FW: 434.4

Chemical Purity: ≥98% (Bicalutamide)

Deuterium

Incorporation: \geq 99% deuterated forms (d₁-d₄); \leq 1% d₀

Supplied as: A solid 4°C Storage: Stability: ≥4 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Bicalutamide- d_{α} is intended for use as an internal standard for the quantification of bicalutamide (Item No. 14250) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Bicalutamide-d₄ is supplied as a solid. A stock solution may be made by dissolving the bicalutamide-d₄ in the solvent of choice. Bicalutamide-d₁ is soluble in organic solvents such as methanol and DMSO, which should be purged with an inert gas.

Description

Bicalutamide is a non-steroidal androgen receptor antagonist that binds the androgen receptor $(K_i = 12.5 \,\mu\text{M}; IC_{50} = 1.2 \,\mu\text{M})$, preventing its activation and subsequent upregulation of androgen responsive genes by androgenic hormones. 1.2 Bicalutamide is frequently used to examine the role of androgen receptor inactivation in the proliferation of prostate cancer cells and has served as a molecular template for the design and structural optimization of more selective androgen receptor modulators for androgen therapy.^{3,4}

References

- 1. Freeman, S.N., Mainwaring, W.I.P., and Furr, B.J.A. A possible explanation for the peripheral selectivity of a novel non-steroidal pure antiandrogen, Casodex (ICI 176,334). Br. J. Cancer 60(5), 664-668 (1989).
- Masiello, D., Cheng, S., Bubley, G.J., et al. Bicalutamide functions as an androgen receptor antagonist by assembly of a transcriptionally inactive receptor. J. Biol. Chem. 277(29), 26321-26326 (2002).
- Gao, W., Kim, J., and Dalton, J.T. Pharmacokinetics and pharmacodynamics of nonsteroidal androgen receptor ligands. Pharm. Res. 23(8), 1641-1658 (2006).
- Yin, D., Perera, M.A., Dalton, J.T., et al. Key structural features of nonsteroidal ligands for binding and activation of the androgen receptor. Mol. Pharmacol. 63(1), 211-223 (2003).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 12/19/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM