



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION

Everolimus-d₄ Item No. 22559

CAS Registry No.: 1338452-54-2
Formal Name: 42-O-(2-hydroxyethyl-1,1,2,2-d₄)-rapamycin
MF: C₅₃H₇₉D₄NO₁₄
FW: 962.3
Chemical Purity: ≥98% (Everolimus)
Deuterium Incorporation: ≥99% deuterated forms (d₁-d₄); ≤1% d₀
Supplied as: A solid
Storage: 4°C
Stability: ≥1 year



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Everolimus-d₄ is intended for use as an internal standard for the quantification of everolimus (Item No. 11597) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Everolimus-d₄ is supplied as a solid. A stock solution may be made by dissolving the everolimus-d₄ in the solvent of choice. Everolimus-d₄ is soluble in organic solvents such as methanol and chloroform, which should be purged with an inert gas.

Description

Everolimus is a hydroxyethyl ether form of rapamycin (Item No. 13346) that inhibits mTOR signaling through both mTORC1 and mTORC2 when added to cells at 20 nM.^{1,2} It is orally available and shows improved pharmacokinetics and pharmacodynamics over rapamycin.² Through its inhibition of mTOR, everolimus inhibits cell proliferation, metabolism, and angiogenesis in certain types of cancer.^{3,4} It also acts as an immunosuppressive agent in the context of organ transplantation.

References

1. Zeng, Z., Sarbassov, D.D., Samudio, I.J., *et al.* Rapamycin derivatives reduce mTORC2 signaling and inhibit AKT activation in AML. *Blood* **109**(8), 3509-3512 (2007).
2. Lebowitz, D., Anak, Ö., Sahmoud, T., *et al.* Development of everolimus, a novel oral mTOR inhibitor, across a spectrum of diseases. *Ann. N.Y. Acad. Sci.* **1291**, 14-32 (2013).
3. Yunokawa, M., Koizumi, F., Kitamura, Y., *et al.* Efficacy of everolimus, a novel mTOR inhibitor, against basal-like triple-negative breast cancer cells. *Cancer Sci.* **103**(9), 1665-1671 (2012).
4. Gurk-Turner, C., Maniatisitkul, W., and Cooper, M. A comprehensive review of everolimus clinical reports: A new mammalian target of rapamycin inhibitor. *Transplantation* **94**(7), 659-668 (2012).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/24/2017

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM