

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Ibrutinib-d₅ Item No. 22561

CAS Registry No.: 1553977-17-5

Formal Name: 1-[(3R)-3-[4-amino-3-[4-(phenoxy-

> 2,3,4,5,6-d₅)phenyl]-1H-pyrazolo[3,4-d] pyrimidin-1-yl]-1-piperidinyl]-2-propen-

 ${\rm C_{25}H_{19}D_5N_6O_2\atop 445.5}$ MF:

FW:


Chemical Purity: Deuterium

Incorporation: \geq 99% deuterated forms (d₁-d₅); \leq 1% d₀

≥98% (Ibrutinib)

A solid Supplied as: -20°C Storage: ≥2 years Stability:

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Ibrutinib-d₅ is intended for use as an internal standard for the quantification of ibrutinib (Item No. 16274) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Ibrutinib- d_5 is supplied as a solid. A stock solution may be made by dissolving the ibrutinib in the solvent of choice. Ibrutinib is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of ibrutinib in DMSO and DMF is approximately 30 mg/ml and approximately 0.25 mg/ml in ethanol.

Description

Ibrutinib is an irreversible inhibitor of Bruton's tyrosine kinase (BTK; $IC_{50} = 0.5$ nM) that selectively blocks B cell activation, promoting apoptosis and preventing homing to the protective tumor microenvironment, at concentrations that do not affect T cell receptor signaling (1,000-fold more potent).^{1,2} It has been reported to inhibit autophosphorylation of BTK (IC_{50} = 11 nM), phosphorylation of PLC γ (IC_{50} = 29 nM), a substrate of BTK, and phosphorylation of ERK (IC_{50} = 13 nM), a further downstream kinase. Formulations containing it have been examined clinically for the treatment of diseases associated with B cell antigen receptor signaling, including mantle cell lymphoma, chronic lymphocytic leukemia, and non-Hodgkin lymphoma. 1,3,4

References

- 1. Honigberg, L. A., Smith, A.M., Sirisawad, M., et al. Proc. Natl. Acad. Sci. USA 107(29), 13075-13080 (2010).
- 2. Herman, S.E.M., Gordon, A.L., Hertlein, E., et al. Blood 117(23), 6287-6296 (2011).
- 3. Leslie, L. A. and Younes, A. Leuk. Lymphoma. 54(11), 2365-2376 (2013).
- 4. Wu, M., Akinleye, A., and Zhu, X. J. Hematol. Oncol. 6, 36 (2013).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/08/2017

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA **PHONE:** [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM